

L I S T

OF THE

BRITISH MARINE INVERTEBRATE FAUNA.

FOR THE DREDGING COMMITTEE OF THE BRITISH ASSOCIATION.

LONDON:

TAYLOR AND FRANCIS, RED LION COURT, FLEET STREET.

1860.

PK-453

L I S T
OF THE
BRITISH MARINE INVERTEBRATE FAUNA.

FOR THE DREDGING COMMITTEE OF THE BRITISH ASSOCIATION.

LONDON:
TAYLOR AND FRANCIS, RED LION COURT, FLEET STREET.
1860.

NOTICE.

THE following lists have been prepared in conformity with the desire of the Committee of the Natural History Section of the British Association for the Advancement of Science, who, at my suggestion, recommended the appointment of a general Dredging Committee, with a liberal grant of money for the carrying out of its objects.

It is intended to place these lists in the hands of the local Dredging Committees and naturalists engaged in researches in the most important districts of the coasts of Great Britain and Ireland, with a request that they may be returned, with notes on the conditions under which each species of the particular district has been found, and memoranda of such additional species as may be obtained. By this means it is hoped to collect local lists of great interest, and materials for a more complete catalogue of the Invertebrate Fauna of the British Seas. In the preparation of the present lists, I have been assisted by Dr. Baird and Mr. S. Woodward, and other members of the Dredging Committee. The catalogue of Mollusca is taken from the work of Messrs. Forbes and Hanley; that of Crustacea has been obligingly furnished by Mr. Spence Bate; of Radiata by Mr. Stewart, of the Royal College of Surgeons; of Sponges by Dr. Bowerbank; of Rhizopoda by Messrs. Rupert Jones and Parker; and to Dr. J. E. Gray I am indebted for permission to extract the list of Anellides from an unpublished work by the late Dr. Johnston, of Berwick-upon-Tweed.

ROBERT McANDREW.

Isleworth House. Feb. 10. 1860.

* * * The nomenclature and arrangement are taken (with a few slight modifications) from the "British Mollusca" of Messrs. Forbes and Hanley.

CEPHALOPODA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
OCTOPUS, <i>Cuvier</i> .					
<i>vulgaris</i> , <i>Lam.</i>					
ELEDONE, <i>Leach</i> .					
<i>octopodia</i> , <i>Penn.</i>					
ROSSIA, <i>Owen</i> .					
<i>Owenii</i> , <i>Ball</i>					
<i>macrosoma</i> , <i>Delle Chiaje</i>					
SEPIOLA, <i>Leach</i> .					
<i>Rondeletii</i> , <i>Leach</i>					
<i>Atlantica</i> , <i>D'Orbigny</i>					
OMMASTREPES, <i>D'Orbigny</i> .					
<i>sagittatus</i> , <i>Lam.</i>					
<i>todarus</i> , <i>Delle Chiaje</i>					
<i>Eblanæ</i> , <i>Ball</i>					
LOLIGO, <i>Lamarck</i> .					
<i>vulgaris</i> , <i>Lam.</i> ? (<i>Forbesi</i> , <i>Stp.</i>)					
<i>media</i> , <i>Linn.</i>					
<i>marmoræ</i> , <i>Verany</i> (<i>media</i> , <i>var.</i>)					
SEPIA, <i>Linnaeus</i> .					
<i>officinalis</i> , <i>Linn.</i>					
<i>elegans</i> , <i>Bl.</i>					
<i>biserialis</i> , <i>De Montfort</i>					

GASTEROPODA.

ORDER PROSOBRANCHIATA.

MUREX, <i>Linnaeus</i> .					
<i>erinaceus</i> , <i>Linn.</i>					
<i>corallinus</i> , <i>Scacchi</i>					
TROPHON, <i>Montfort</i> .					
<i>clathratus</i> , <i>Linn.</i>					
<i>muricatus</i> , <i>Mont.</i>					
<i>Barviensis</i> , <i>Johnston</i>					
FUSUS, <i>Lamarck</i> .					
<i>gracilis</i> , <i>Da Costa</i>					
<i>propinquus</i> , <i>Alder</i>					
<i>Berniciensis</i> , <i>King</i>					
<i>antiquus</i> , <i>Linn.</i>					
<i>Norvegicus</i> , <i>Chemn.</i>					
<i>Turtoni</i> , <i>Bean</i>					
BUCCINUM, <i>Linnaeus</i> .					
<i>undatum</i> , <i>Linn.</i>					
<i>Dalei</i> , <i>J. Sowerby</i>					
<i>Humphresianum</i> , <i>Bennett</i>					
<i>fusiforme</i> , <i>Broderip</i>					

GASTEROPODA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
NASSA, <i>Lamarck</i> .					
<i>reticulata</i> , <i>Linn.</i>					
<i>pygmæa</i> , <i>Lamarck</i>					
<i>incrassata</i> , <i>Müller</i>					
PURPURA, <i>Lamarck</i> .					
<i>lapillus</i> , <i>Linn.</i>					
MANGELIA, <i>Leach</i> .					
<i>attenuata</i> , <i>Mont.</i>					
<i>costata</i> , <i>Pennant</i>					
<i>brachystoma</i> , <i>Philippi</i>					
<i>gracilis</i> , <i>Mont.</i>					
<i>Leufroyi</i> , <i>Michaud</i>					
<i>linearis</i> , <i>Mont.</i>					
<i>nana</i> , <i>Lovén</i>					
<i>nebula</i> , <i>Mont.</i>					
<i>purpurea</i> , <i>Mont.</i>					
<i>rufa</i> , <i>Mont.</i>					
<i>septangularis</i> , <i>Mont.</i>					
<i>striolata</i> , <i>Scacchi</i>					
<i>teres</i> , <i>Forbes</i>					
<i>Trevelliana</i> , <i>Turton</i>					
<i>turricula</i> , <i>Mont.</i>					
LACHESIS, <i>Risso</i> .					
<i>minima</i> , <i>Mont.</i>					
MARGINELLA, <i>Lamarck</i> .					
<i>lævis</i> , <i>Donovan</i>					
OVULA, <i>Lamarck</i> .					
<i>patula</i> , <i>Pennant</i>					
? <i>acuminata</i> , <i>Bruguère</i>					
CYPRÆA, <i>Linnaeus</i> .					
<i>Europæa</i> , <i>Mont.</i>					
NATICA, <i>Lamarck</i> .					
<i>monilifera</i> , <i>Lamarck</i>					
<i>nitida</i> , <i>Donovan</i>					
<i>sordida</i> , <i>Philippi</i>					
<i>Montagui</i> , <i>Forbes</i>					
<i>helicoides</i> , <i>Johnston</i>					
<i>pusilla</i> , <i>Say</i>					
<i>Kingii</i> , <i>Forbes</i>					
LAMELLARIA, <i>Montagu</i> .					
<i>perspicua</i> , <i>Linn.</i>					
<i>tentaculata</i> , <i>Mont.</i>					
VELUTINA, <i>Fleming</i> .					
<i>flexilis</i> , <i>Mont.</i>					
<i>lævigata</i> , <i>Linn.</i>					
? OTINA, <i>Gray</i> .					
<i>otis</i> , <i>Turton</i>					
TRICHOtropis, <i>Broderip</i> .					
<i>borealis</i> , <i>Brod.</i>					

GASTEROPODA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
<i>CERITHIOPSIS, Forbes & Hanley.</i>					
<i>tubercularis, Mont.</i>					
<i>ODOSTOMIA, Fleming.</i>					
<i>acuta, Jeffreys</i>					
<i>alba, Jeffreys</i>					
<i>conoidea, Brocchi</i>					
<i>conspicua, Alder</i>					
<i>cylindrica, Alder</i>					
<i>decussata, Mont.</i>					
<i>dolioliformis, Jeffreys</i>					
<i>dubia, Jeffreys</i>					
<i>eulimoides, Hanley</i>					
<i>excavata, Philippi</i>					
<i>glabrata, Mühlfeldt</i>					
<i>Gulsonæ, Clark</i>					
<i>insculpta, Mont.</i>					
<i>interstincta, Mont.</i>					
<i>minuta, Jeffreys</i>					
<i>nitida, Alder</i>					
<i>obliqua, Alder</i>					
<i>pallida, Mont.</i>					
<i>plicata, Mont.</i>					
<i>rissoides, Hanley</i>					
<i>spiralis, Mont.</i>					
<i>striolata, Alder</i>					
<i>truncatula, Jeffreys</i>					
<i>unidentata, Mont.</i>					
<i>Warrenii, Thompson</i>					
<i>EULIMELLA, Forbes.</i>					
<i>acicula, Philippi</i>					
<i>affinis, Philippi</i>					
<i>clavula, Lovén</i>					
<i>Scillæ, Scacchi</i>					
<i>CHEMNITZIA, D'Orbigny.</i>					
<i>clathrata, Jeffreys</i>					
<i>elegantissima, Mont.</i>					
<i>fenestrata, Jeffreys</i>					
<i>formosa, Jeffreys</i>					
<i>fulvocincta, Thomps.</i>					
<i>indistincta, Mont.</i>					
<i>rufa, Philippi</i>					
<i>rufescens, Forbes</i>					
<i>scalaris, Philippi</i>					
<i>eximia, Jeffreys</i>					
<i>EULIMA, Risso.</i>					
<i>polita, Linn.</i>					
<i>distorta, Deshayes</i>					
<i>subulata, Donovan</i>					
<i>bilineata, Alder</i>					

GASTEROPODA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
STYLINA, <i>Fleming</i> .					
<i>Turtoni</i> , <i>Broderip</i>					
CERITHIUM, <i>Bruguère</i> .					
<i>metula</i> , <i>Lovén</i>					
<i>reticulatum</i> , <i>Da Costa</i>					
<i>adversum</i> , <i>Mont.</i>					
APORRHÆIS, <i>Aldrovandus</i> .					
<i>pes-carbonis</i> , <i>Brongniart</i>					
<i>pes-pelecani</i> , <i>Linn.</i>					
TURRITELLA, <i>Lamarck</i> .					
<i>communis</i> , <i>Risso</i>					
ACLIS, <i>Lovén</i> .					
<i>ascaris</i> , <i>Turton</i>					
<i>supranitida</i> , <i>S. Wood</i>					
? <i>unica</i> , <i>Mont.</i>					
<i>nitidissima</i> , <i>Mont.</i>					
CLECUM, <i>Fleming</i> .					
<i>trachea</i> , <i>Mont.</i>					
<i>glabrum</i> , <i>Mont.</i>					
SCALARIA, <i>Lamarck</i> .					
<i>Turtoni</i> , <i>Turton</i>					
<i>communis</i> , <i>Lamarck</i>					
<i>clathratula</i> , <i>Mont.</i>					
<i>Grœnlandica</i> , <i>Chemnitz</i>					
<i>Trevelyana</i> , <i>Leach</i>					
SKENEIA, <i>Fleming</i> .					
? <i>costulata</i> , <i>Müller</i>					
? <i>lævis</i> , <i>Philippi</i>					
<i>planorbis</i> , <i>Fabr.</i>					
? <i>nitidissima</i> , <i>Adams</i>					
? <i>rota</i> , <i>Forbes</i>					
TRUNCATELLA, <i>Risso</i> .					
<i>Montagui</i> , <i>Lowe</i>					
JEFFREYSIA, <i>Alder</i> .					
<i>opalina</i> , <i>Jeffreys</i>					
<i>diaphana</i> , <i>Alder</i>					
<i>globularis</i> , <i>Jeffreys</i>					
RISSEA, <i>Frémenville</i> .					
<i>abyssicola</i> , <i>Forbes</i>					
<i>anatina</i> , <i>Drap.</i>					
<i>Beanii</i> , <i>Hanley</i>					
<i>calathus</i> , <i>Forbes</i>					
<i>cingillus</i> , <i>Mont.</i>					
<i>costata</i> , <i>Adams</i>					
<i>costulata</i> , <i>Risso</i>					
<i>crenulata</i> , <i>Michaud</i>					
<i>fulgida</i> , <i>Adams</i>					
<i>inconspicua</i> , <i>Alder</i>					
<i>labiosa</i> , <i>Mont.</i>					

GASTEROPODA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
RISSEO , <i>Frémeville</i> .					
<i>lactea</i> , <i>Michaud</i>					
<i>littorea</i> , <i>Delle Chiaje</i>					
<i>muriatica</i> , <i>Lam.</i>					
<i>parva</i> , <i>Da Costa</i>					
<i>proxima</i> , <i>Alder</i>					
<i>pulcherrima</i> , <i>Jeffreys</i>					
<i>punctura</i> , <i>Mont.</i>					
<i>rubra</i> , <i>Adams</i>					
<i>rufilabrum</i> , <i>Alder</i>					
<i>sculpta</i> , <i>Philippi</i>					
<i>semistriata</i> , <i>Mont.</i>					
<i>soluta</i> , <i>Philippi</i>					
<i>striata</i> , <i>Mont.</i>					
<i>striatula</i> , <i>Mont.</i>					
<i>ulvæ</i> , <i>Pennant</i>					
<i>ventrosa</i> , <i>Mont.</i>					
<i>vitrea</i> , <i>Mont.</i>					
<i>Zetlandica</i> , <i>Mont.</i>					
ASSIMINEA , <i>Leach</i> .					
<i>Grayana</i> , <i>Leach</i>					
LACUNA , <i>Turton</i> .					
<i>crassior</i> , <i>Mont.</i>					
<i>vincta</i> , <i>Mont.</i>					
<i>puteolus</i> , <i>Turton</i>					
<i>pallidula</i> , <i>Da Costa</i>					
LITORINA , <i>Férussac</i> .					
<i>fabalis</i> , <i>Turton</i>					
<i>litoralis</i> , <i>Linn.</i>					
<i>litorea</i> , <i>Linn.</i>					
<i>neritoides</i> , <i>Linn.</i>					
<i>tenebrosa</i> , <i>Mont.</i>					
<i>palliata</i> , <i>Say</i>					
<i>patula</i> , <i>Jeffreys</i>					
<i>rudis</i> , <i>Donovan</i>					
<i>saxatilis</i> , <i>Johnston</i>					
ADEORBIS , <i>S. Wood</i> .					
<i>subcarinata</i> , <i>Mont.</i>					
<i>divisa</i> , <i>Fleming</i>					
TROCHUS , <i>Linn.</i>					
<i>alabastrum</i> , <i>Beck</i>					
<i>cinerarius</i> , <i>Linn.</i>					
<i>conulus</i> , <i>Linn.</i>					
<i>exiguus</i> , <i>Pulteney</i>					
<i>granulatus</i> , <i>Born</i>					
<i>crassus</i> , <i>Pulteney</i> (<i>lineatus</i> , <i>Da Costa</i>)					
<i>magus</i> , <i>Linn.</i>					
<i>millegranus</i> , <i>Philippi</i>					

GASTEROPODA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
<i>TROCHUS, Linn.</i>					
<i>Montagui, Gray</i>					
<i>striatus, Linn.</i>					
<i>tumidus, Mont.</i>					
<i>umbilicatus, Mont.</i>					
<i>lineatus, Da Costa</i>					
<i>zizyphinus, Linn.</i>					
<i>MARGARITA, Leach.</i>					
<i>undulata, Sowerby</i>					
<i>helicina, Fabr</i>					
<i>pusilla, Jeffreys</i>					
<i>Cutleriana, Clark</i>					
<i>PHASIANELLA, Lamarck.</i>					
<i>pullus, Linn.</i>					
<i>LANTHINA, Lamarck.</i>					
<i>exigua, Lamarck</i>					
<i>communis, Lamarck.</i>					
<i>pallida, Harvey</i>					
<i>SCISSURELLA, D'Orbigny.</i>					
<i>crispata, Fleming.</i>					
<i>HALIOTIS, Linn.</i>					
<i>tuberculata, Linn.</i>					
<i>EMARGINULA, Lamarck.</i>					
<i>reticulata, J. Sow.</i>					
<i>rosea, Bell</i>					
<i>crassa, J. Sow.</i>					
<i>PUNCTURELLA, Lowe.</i>					
<i>Noachina, Linn.</i>					
<i>FISSURELLA, Lamarck.</i>					
<i>reticulata, Donovan</i>					
<i>PILEOPSIS, Lamarck.</i>					
<i>Hungaricus, Linn.</i>					
<i>CALYPTREÆ, Lamarck.</i>					
<i>Sincensis, Linn.</i>					
<i>ACMÆA, Eschscholtz.</i>					
<i>testudinalis, Müller</i>					
<i>virginea, Müller</i>					
<i>PATELLA, Linnæus.</i>					
<i>vulgata, Linn.</i>					
<i>athletica, Bean</i>					
<i>pellucida, Linn.</i>					
<i>lævis, Pennant</i>					
<i>PILIDIUM, Forbes.</i>					
<i>fulvum, Müller</i>					
<i>PROPILIDIUM, Forbes. (LEPETA, Gr.)</i>					
<i>ancyloide, Forbes</i>					
<i>DENTALIUM, Linnæus.</i>					
<i>entale, Linn.</i>					
<i>Tarentinum, Lam.</i>					

GASTEROPODA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
CHITON, <i>Linnaeus</i> .	fathoms.	fathoms.			
<i>fascicularis</i> , <i>Linn.</i>					
<i>discrepans</i> , <i>Brown</i>					
<i>Hanleyi</i> , <i>Bean</i>					
<i>ruber</i> , <i>Linn.</i>					
<i>cinereus</i> , <i>Linn</i>					
<i>albus</i> , <i>Linn.</i>					
<i>asellus</i> , <i>Chemn.</i>					
<i>cancellatus</i> , <i>Sow.</i>					
<i>laevis</i> , <i>Pennant</i>					
<i>marmoreus</i> , <i>O. Fabr.</i>					

ORDER OPISTHOBRANCHIATA.

TORNATELLA, <i>Lamarck</i> .					
<i>fasciata</i> , <i>Linn.</i>					
BULLA, <i>Lamarck</i> .					
<i>hydatis</i> , <i>Linn.</i>					
<i>Cranchii</i> , <i>Leach</i>					
AKEREA, <i>Müller</i> .					
<i>bullata</i> , <i>Linn.</i>					
CYLICHNA, <i>Lovén</i> .					
<i>cylindracea</i> , <i>Pennant</i>					
<i>conulus</i> , <i>Desh.</i>					
<i>mamillata</i> , <i>Philippi</i>					
<i>nitidula</i> , <i>Lovén</i>					
<i>obtusa</i> , <i>Mont.</i>					
<i>strigella</i> , <i>Lovén</i>					
<i>truncata</i> , <i>Adams</i>					
<i>umbilicata</i> , <i>Mont.</i>					
AMPHISPHYRA, <i>Lovén</i> .					
<i>hyalina</i> , <i>Turton</i>					
SCAPHANDER, <i>Montfort</i> .					
<i>lignarius</i> , <i>Linn.</i>					
BULLÆA, <i>Lamarck</i> .					
<i>aperta</i> , <i>Linn.</i>					
<i>quadrata</i> , <i>S. Wood</i>					
<i>scabra</i> , <i>Müller</i>					
<i>catena</i> , <i>Mont.</i>					
<i>punctata</i> , <i>Clark</i>					
<i>pruinosa</i> , <i>Clark</i>					
APLYSIA, <i>Gmelin</i> .					
<i>hybrida</i> , <i>Sow.</i>					
PLEUROBRANCHUS, <i>Cuvier</i> .					
<i>plumula</i> , <i>Mont.</i>					
<i>membranaceus</i> , <i>Mont.</i>					
DIPHYLLIDIA, <i>Cuvier</i> .					
<i>lineata</i> , <i>Otto</i>					

7

GASTEROPODA.

ORDER NUDIBRANCHIATA.

(From the Monograph of Messrs. Alder and Hancock, 1856.)

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
<i>DORIS, Linn.</i>					
tuberculata, <i>Cuv.</i>					
flammea, <i>A. & H.</i>					
Zetlandica, <i>A. & H.</i>					
millegrana, <i>A. & H.</i>					
Johnstoni, <i>A. & H.</i>					
planata, <i>A. & H.</i>					
coccinea, <i>Forbes</i>					
repanda, <i>A. & H.</i>					
aspera, <i>A. & H.</i>					
proxima, <i>A. & H.</i>					
muricata, <i>Müll.</i>					
Ulidiana, <i>Thomps.</i>					
diaphana, <i>A. & H.</i>					
oblonga, <i>A. & H.</i>					
bilamellata, <i>L.</i>					
depressa, <i>A. & H.</i>					
inconspicua, <i>A. & H.</i>					
pusilla, <i>A. & H.</i>					
sparsa, <i>A. & H.</i>					
pilosa, <i>Müll.</i>					
subquadrata, <i>A. & H.</i>					
<i>GONIODORIS, Forbes.</i>					
nodosa, <i>Mont.</i>					
castanea, <i>A. & H.</i>					
<i>TRIOPA, Johnston.</i>					
claviger, <i>Müll.</i>					
<i>ÆGIRUS, Lovén.</i>					
punctilucens, <i>D'Orb.</i>					
<i>THECACERA, Fleming.</i>					
pennigera, <i>Mont.</i>					
virescens, <i>A. & H.</i>					
capitata, <i>A. & H.</i>					
<i>POLYCERA, Cuvier.</i>					
quadrilineata, <i>Müll.</i>					
ocellata, <i>A. & H.</i>					
Lessonii, <i>D'Orb.</i>					
<i>ANCUA, Lovén.</i>					
cristata, <i>Alder</i>					
<i>IDALIA, Leuckart.</i>					
elegans, <i>Leuck.</i>					
Leachii, <i>A. & H.</i>					
aspersa, <i>A. & H.</i>					

GASTEROPODA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
<i>IDALIA, Leuckart.</i>					
<i>inaequalis, Forbes</i>					
<i>pulchella, A. & H.</i>					
<i>quadricornis, Mont.</i>					
<i>TRITONIA, Cuvier.</i>					
<i>Homborgii, Cuv.</i>					
<i>alba, A. & H.</i>					
<i>plebeia, Johnston</i>					
<i>lineata, A. & H.</i>					
<i>SCYLLÆA, Linn.</i>					
<i>pelagica, Linn.</i>					
<i>LOMANOTUS, Verany.</i>					
<i>marmoratus, A. & H.</i>					
<i>flavidus, A. & H.</i>					
<i>DENDRONOTUS, A. & H.</i>					
<i>arborescens, Müll.</i>					
<i>DOTO, Oken.</i>					
<i>fragilis, Forbes</i>					
<i>pinnatifida, Mont.</i>					
<i>coronata, Müll.</i>					
<i>ÆOLIS, Cuvier.</i>					
<i>papillosa, Linn.</i>					
<i>glauca, A. & H.</i>					
<i>Alderi, Cocks</i>					
<i>coronata, Forbes</i>					
<i>Drummondi, Thomps.</i>					
<i>punctata, A. & H.</i>					
<i>elegans, A. & H.</i>					
<i>rufibranchialis, Johnst.</i>					
<i>lineata, Lov.</i>					
<i>smaragdina, A. & H.</i>					
<i>gracilis, A. & H.</i>					
<i>pellucida, A. & H.</i>					
<i>Landsburgii, A. & H.</i>					
<i>alba, A. & H.</i>					
<i>carnea, A. & H.</i>					
<i>glaucoides, A. & H.</i>					
<i>Peachii, A. & H.</i>					
<i>nana, A. & H.</i>					
<i>stipata, A. & H.</i>					
<i>angulata, A. & H.</i>					
<i>inornata, A. & H.</i>					
<i>concinna, A. & H.</i>					
<i>olivacea, A. & H.</i>					
<i>aurantiaca, A. & H.</i>					
<i>pustulata, A. & H.</i>					
<i>Couchii, Cocks</i>					
<i>amœna, A. & H.</i>					
<i>Northumbria, A. & H.</i>					

GASTEROPODA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
<i>ÆOLIS, Cuvier.</i>					
arenicola, Forbes					
Glottensis, A. & H.					
viridis, Forbes					
purpurascens, Flem.					
cingulata, A. & H.					
vittata, A. & H.					
cærulea, Mont.					
pieta, A. & H.					
tricolor, Forbes.					
amethystina, A. & H.					
Farrani, A. & H.					
exigua, A. & H.					
despecta, Johnst.					
<i>EMBLETONIA, A. & H.</i>					
pulchra, A. & H.					
minuta, F. & G.					
pallida, A. & H.					
<i>FIONA, A. & H.</i>					
nobilis, A. & H.					
<i>HERMEA, Lovén.</i>					
bifida, Mont.					
dendritica, A. & H.					
<i>ALDERIA, Allman.</i>					
modesta, Lovén					
<i>PROCTONOTUS, A. & H.</i>					
mucroniferus, A. & H.					
<i>ANTIOPA, A. & H.</i>					
cristata, Del. Ch.					
hyalina, A. & H.					

PTEROPODA.

<i>HYALEA, Lamarck.</i>					
trispinosa, Lesueur					
<i>SPIRIALIS, Eyloux & Souleyet.</i>					
Flemingii, Forbes					
Jeffreysii, Forbes					
MacAndrei, Forbes					
<i>CLIO, Müller.</i>					
borealis, Linn.					

LAMELLIBRANCHIATA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
OSTREA, <i>Linnaeus</i> .					
<i>edulis</i> , <i>Linn.</i>					
ANOMIA, <i>Linnaeus</i> .					
<i>aculeata</i> , <i>Müller</i>					
<i>ephippium</i> , <i>Linn.</i>					
<i>striata</i> , <i>Lovén</i>					
<i>patelliformis</i> , <i>Linn.</i>					
PECTEN, <i>O. F. Müller</i> .					
<i>Danicus</i> , <i>Chemnitz</i>					
<i>maximus</i> , <i>Linn.</i>					
<i>niveus</i> , <i>Macgillivray</i>					
<i>opercularis</i> , <i>Linn.</i>					
<i>pusio</i> , <i>Pennant</i>					
<i>similis</i> , <i>Laskey</i>					
<i>tigrinus</i> , <i>Müller</i>					
<i>varius</i> , <i>Linn.</i>					
<i>striatus</i> , <i>Müller</i>					
<i>furtivus</i> , <i>Lovén</i>					
LIMA, <i>Bruguère</i> .					
<i>hians</i> , <i>Gmelin</i>					
<i>Loscombii</i> , <i>Sowerby</i>					
<i>subauriculata</i> , <i>Mont.</i>					
AVICULA, <i>Bruguère</i> .					
<i>Tarentina</i> , <i>Lam.</i>					
PINNA, <i>Linnaeus</i> .					
<i>pectinata</i> , <i>Linn.</i>					
MYTILUS, <i>Linnaeus</i> .					
<i>edulis</i> , <i>Linn.</i>					
MODIOLA, <i>Lamarck</i> .					
<i>barbata</i> , <i>Linn.</i>					
<i>modiolus</i> , <i>Linn.</i>					
<i>phaseolina</i> , <i>Philippi</i>					
<i>tulipa</i> , <i>Lam.</i>					
CRENELLA, <i>Brown</i> .					
<i>costulata</i> , <i>Risso</i>					
<i>decussata</i> , <i>Mont.</i>					
<i>discors</i> , <i>Linn.</i>					
<i>nigra</i> , <i>Gray</i>					
<i>marmorata</i> , <i>Forbes</i>					
<i>rhombea</i> , <i>Berkeley</i>					
ARCA, <i>Linnaeus</i> .					
<i>lactea</i> , <i>Linn.</i>					
<i>raridentata</i> , <i>S. Wood</i>					
<i>tetragona</i> , <i>Poli</i>					
PECTUNCULUS, <i>Lamarck</i> .					
<i>glycimeris</i> , <i>Linn.</i>					

LAMELLIBRANCHIATA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
<i>NUCULA, Lamarck.</i>					
<i>decussata, Sow.</i>					
<i>nitida, Sow.</i>					
<i>nucleus, Linn.</i>					
<i>radiata, Hanley</i>					
<i>tenuis, Mont.</i>					
<i>LEDA, Schumacher.</i>					
<i>caudata, Don.</i>					
<i>pygmæa, Münster</i>					
<i>CARDIUM, Linnaeus.</i>					
<i>aculeatum, Linn.</i>					
<i>echinatum, Linn.</i>					
<i>edule, Linn.</i>					
<i>fasciatum, Mont.</i>					
<i>nodosum, Turton</i>					
<i>Norvegicum, Speng.</i>					
<i>pygmæum, Don.</i>					
<i>rusticum, Linn.</i>					
<i>Succicum, Reeve</i>					
<i>LUCINA, Bruguière.</i>					
<i>borealis, Linn.</i>					
<i>divaricata, Linn.</i>					
<i>ferruginosa, Forbes</i>					
<i>flexuosa, Mont.</i>					
<i>leucoma, Turton</i>					
<i>spinifera, Mont.</i>					
<i>DIPLODONTA, Bronn.</i>					
<i>rotundata, Mont.</i>					
<i>KELLIA, Turton.</i>					
<i>suborbicularis, Mont.</i>					
<i>rubra, Mont.</i>					
<i>TURTONIA, Hanley.</i>					
<i>minuta, O. Fabr.</i>					
<i>MONTACUTA, Turton.</i>					
<i>bidentata, Mont.</i>					
<i>ferruginosa, Mont.</i>					
<i>substriata, Mont.</i>					
<i>LEPTON, Turton.</i>					
<i>Clarkiæ, Clark</i>					
<i>nitidum, Turton</i>					
<i>squamosum, Mont.</i>					
<i>GALEOMMA, Turton.</i>					
<i>Turtoni, Sow.</i>					
<i>CYPRINA, Lamarck.</i>					
<i>Islandica, Linn.</i>					
<i>CIRCE, Schumacher.</i>					
<i>minima, Mont.</i>					
<i>ASTARTE, Sowerby.</i>					
<i>arctica, Gray</i>					

LAMELLIBRANCHIATA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
ASTARTE, Sowerby.					
compressa, <i>Mont.</i>					
crebricostata, <i>Forbes</i>					
elliptica, <i>Brown</i>					
sulcata, <i>Da Costa</i>					
triangularis, <i>Mont.</i>					
ISOCARDIA, Lamarck.					
cor, <i>Linn.</i>					
VENUS, Linnæus.					
casina, <i>Linn.</i>					
fasciata, <i>Don.</i>					
ovata, <i>Pennant.</i>					
striatula, <i>Don.</i>					
verrucosa, <i>Linn.</i>					
CYTHEREA, Lamarck.					
chione, <i>Linn.</i>					
ARTEMIS, Poli.					
exoleta, <i>Linn.</i>					
lineta, <i>Pult.</i>					
LUCINOPSIS, Forbes.					
undata, <i>Penn.</i>					
TAPES, Mühlfeldt.					
aurea, <i>Gmelin</i>					
decussata, <i>Linn.</i>					
pullastra, <i>Wood</i>					
virginica, <i>Linn.</i>					
VENERUPIS, Lamarck.					
irus, <i>Linn.</i>					
PETRICOLA, Lamarck.					
lithophaga, <i>Retzius</i>					
MACTRA, Linnæus.					
elliptica, <i>Brown</i>					
helvacea, <i>Chemnitz</i>					
solida, <i>Linn.</i>					
stultorum, <i>Linn.</i>					
subtruncata, <i>Da Costa</i>					
truncata, <i>Mont.</i>					
LUTRARIA, Lamarck.					
elliptica, <i>Linn.</i>					
oblonga, <i>Chemn.</i>					
TELLINA, Linnæus.					
balaustina, <i>Linn.</i>					
crassa, <i>Penn.</i>					
donacina, <i>Linn.</i>					
fabula, <i>Gronov.</i>					
incarnata, <i>Linn.</i>					
proxima, <i>Brown</i>					
pygmaea, <i>Philippi</i>					
solidula, <i>Pult.</i>					

LAMELLIBRANCHIATA.

Species.	Range.	Found living at	Ground.	Frequency	Observations.
	fathoms.	fathoms.			
TELLINA, <i>Linnaeus</i> .					
<i>tenuis</i> , <i>Da Costa</i>					
GASTRANA, <i>Sch.</i> (DIODONTA, <i>F. & H.</i>)					
<i>fragilis</i> , <i>Linn.</i>					
PSAMMOBIA, <i>Lamarck</i> .					
<i>costulata</i> , <i>Turt.</i>					
<i>Ferroensis</i> , <i>Chemn.</i>					
<i>tellinella</i> , <i>Lam.</i>					
<i>vespertina</i> , <i>Chemn.</i>					
SYNDOSMYA, <i>Recluz</i> .					
<i>alba</i> , <i>Wood</i>					
<i>intermedia</i> , <i>Thomps.</i>					
<i>prismatica</i> , <i>Mont.</i>					
<i>tenuis</i> , <i>Mont.</i>					
SCROBICULARIA, <i>Schumacher</i> .					
<i>piperata</i> , <i>Gmelin</i>					
ERVILIA, <i>Turton</i> .					
<i>castanea</i> , <i>Mont.</i>					
DONAX, <i>Linnaeus</i> .					
<i>anatinus</i> , <i>Lam.</i>					
<i>politus</i> , <i>Poli</i>					
SOLEN, <i>Linnaeus</i> .					
<i>ensis</i> , <i>Linn.</i>					
<i>marginatus</i> , <i>Pult.</i>					
<i>pellucidus</i> , <i>Penn.</i>					
<i>siliqua</i> , <i>Linn.</i>					
CERATISOLEN, <i>Forbes</i> .					
<i>legumen</i> , <i>Linn.</i>					
SOLECURTUS, <i>Blainville</i> .					
<i>candidus</i> , <i>Renieri</i>					
<i>coarctatus</i> , <i>Gmelin</i>					
MYA, <i>Linnaeus</i> .					
<i>arenaria</i> , <i>Linn.</i>					
<i>truncata</i> , <i>Linn.</i>					
CORBULA, <i>Bruquière</i> .					
<i>nucleus</i> , <i>Lam.</i>					
<i>ovata</i> , <i>Forbes</i>					
<i>rosea</i> , <i>Brown</i>					
SPHENIA, <i>Turton</i> .					
<i>Binghami</i> , <i>Turton</i>					
NEÆRA, <i>Gray</i> .					
<i>abbreviata</i> , <i>Forbes</i>					
<i>costellata</i> , <i>Desh.</i>					
<i>cuspidata</i> , <i>Oliv.</i>					
POROMYA, <i>Forbes</i> . (=THETIS, <i>Sby.</i>)					
<i>granulata</i> , <i>Nyst</i>					
PANOPÆA, <i>Menard de la Groye</i> .					
<i>Norvegica</i> , <i>Speng.</i>					

LAMELLIBRANCHIATA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
SAXICAVA, <i>Bellevue</i> .	fathoms.	fathoms.			
<i>arctica</i> , <i>Linn.</i>					
<i>rugosa</i> , <i>Linn.</i>					
COCHLODESMA, <i>Leach.</i> (= <i>PERIPLOMA</i> , <i>Sch.</i>)					
<i>prætenue</i> , <i>Pult.</i>					
THRACIA, <i>Leach.</i>					
<i>convexa</i> , <i>Wood</i>					
<i>distorta</i> , <i>Mont.</i>					
<i>phaseolina</i> , <i>Lam.</i>					
<i>pubescens</i> , <i>Pult.</i>					
<i>villosiuscula</i> , <i>Macgill.</i>					
LYONSIA, <i>Turton.</i>					
<i>Norvegica</i> , <i>Chemn.</i>					
PANDORA, <i>Bruquière.</i>					
<i>obtusa</i> , <i>Leach</i>					
<i>rostrata</i> , <i>Lam.</i>					
GASTROCHÆNA, <i>Spengler.</i>					
<i>modiolina</i> , <i>Lam.</i>					
PHOLAS, <i>Linnaeus.</i>					
<i>candida</i> , <i>Linn.</i>					
<i>crispata</i> , <i>Linn.</i>					
<i>dactylus</i> , <i>Linn.</i>					
<i>parva</i> , <i>Penn.</i>					
<i>striata</i> , <i>Linn.</i>					
PHOLADIDEA, <i>Turton.</i>					
<i>lamellata</i> , <i>Turton</i>					
<i>papyracea</i> , <i>Solander.</i>					
XYLOPHAGA, <i>Turton.</i>					
<i>dorsalis</i> , <i>Turton</i>					
TEREDO, <i>Adanson.</i>					
<i>bipennata</i> , <i>Turton</i>					
<i>malleolus</i> , <i>Turton</i>					
<i>megotara</i> , <i>Hanley</i>					
<i>navalis</i> , <i>Linn.</i>					
<i>Norvegica</i> , <i>Speng.</i>					
<i>palmulata</i> , <i>Lam.</i>					

BRACHIOPODA.

CRANIA, <i>Retzius.</i>					
<i>anomala</i> , <i>Müller</i>					
RHYNCHONELLA, <i>Fischer.</i>					
<i>psittacea</i> , <i>Chemn.</i>					
TEREBRATULA, <i>Bruquière.</i>					
<i>caput serpentis</i> , <i>Linn.</i>					
<i>cranium</i> , <i>Müller</i>					
<i>capsula</i> , <i>Jeffreys</i>					
ARGIOPE, <i>Deslongchamps.</i>					
<i>cistellula</i> , <i>Searles Wood</i>					

TUNICATA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			3
APLIDIUM, <i>Savigny</i> .					
<i>ficus</i> , <i>Linn.</i>					
<i>fallax</i> , <i>Johnst.</i>					
<i>nutans</i> , <i>Johnst.</i>					
SIDNYUM, <i>Savigny</i> .					
<i>turbinatum</i> , <i>Savig.</i>					
POLYCLINUM, <i>Savigny</i> .					
<i>aurantium</i> , <i>M.-Edw.</i>					
AMOUROUCIUM, <i>M.-Edw.</i>					
<i>proliferum</i> , <i>M.-Edw.</i>					
<i>Nordmanni</i> , <i>M.-Edw.</i>					
<i>Argus</i> , <i>M.-Edw.</i>					
LEPTOCLINUM, <i>M.-Edw.</i>					
<i>maculosum</i> , <i>M.-Edw.</i>					
<i>asperum</i> , <i>M.-Edw.</i>					
<i>aureum</i> , <i>M.-Edw.</i>					
<i>gelatinosum</i> , <i>M.-Edw.</i>					
<i>Listerianum</i> , <i>M.-Edw.</i>					
<i>punctatum</i> , <i>Forbes</i>					
DISTOMA, <i>Gaertner</i> .					
<i>rubrum</i> , <i>Savig.</i>					
<i>variolosum</i> , <i>Gaertner</i>					
BOTRYLLUS, <i>Gaertner</i> .					
<i>Schlosseri</i> , <i>Pallas</i>					
<i>polycyclus</i> , <i>Savig.</i>					
<i>gemmeus</i> , <i>Savig.</i>					
<i>violaceus</i> , <i>M.-Edw.</i>					
<i>smaragdus</i> , <i>M.-Edw.</i>					
<i>virescens</i> , <i>A. & H.</i>					
<i>bivittatus</i> , <i>M.-Edw.</i>					
<i>rubens</i> , <i>A. & H.</i>					
<i>castaneus</i> , <i>A. & H.</i>					
BOTRYLLOIDES, <i>M.-Edw.</i>					
<i>Leachii</i> , <i>Savig.</i>					
<i>ramulosa</i> , <i>A. & H.</i>					
<i>albicans</i> , <i>M.-Edw.</i>					
<i>radiata</i> , <i>A. & H.</i>					
<i>rotifera</i> , <i>M.-Edw.</i>					
<i>rubra</i> , <i>M.-Edw.</i>					
CLAVELINA, <i>Savigny</i> .					
<i>lepadiformis</i> , <i>O. F. Müller</i>					
PEROPHORA, <i>Wiegmann</i> .					
<i>Listeri</i> , <i>Wiegmann</i>					
SYNTETHYS, <i>Forbes & Goodsir</i> .					
<i>Hebridicus</i> , <i>F. & G.</i>					

TUNICATA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
ASCIDIA, Baster.					
intestinalis, <i>Linn.</i>					
canina, <i>O. F. Müll.</i>					
venosa, <i>O. F. Müll.</i>					
mentula, <i>O. F. Müll.</i>					
arachnoïdea, <i>E. Forbes</i>					
scabra, <i>O. F. Müll.</i>					
virginea, <i>O. F. Müll.</i>					
parallélogramma, <i>O. F. Müll.</i>					
prunum, <i>Müll.?</i>					
orbicularis, <i>Müll.</i>					
depressa, <i>A. & H.</i>					
aspersa, <i>Müll.</i>					
vitrea, <i>Van Beneden</i>					
conchilega, <i>O. F. Müll.</i>					
echinata, <i>Linn.</i>					
sordida, <i>A. & H.</i>					
albida, <i>A. & H.</i>					
elliptica, <i>A. & H.</i>					
pellucida, <i>A. & H.</i>					
MOLGULA, E. Forbes.					
oculata, <i>E. Forbes</i>					
arenosa, <i>A. & H.</i>					
CYNTHIA, Savigny.					
microcosmus, <i>Savig.</i>					
claudicans, <i>Savig.</i>					
tuberosa, <i>Macgillivray</i>					
quadrangularis, <i>E. Forbes</i>					
informis, <i>E. Forbes</i>					
tessellata, <i>E. Forbes</i>					
limacina, <i>E. Forbes</i>					
morus, <i>E. Forbes</i>					
rustica, <i>Linn.</i>					
grossularia, <i>Van Beneden</i>					
ampulla, <i>Brug.</i>					
mamillaris, <i>Pallas</i>					
aggregata, <i>Rathke</i>					
coriacea, <i>A. & H.</i>					
PELONAIIA, Forbes & Goodsir.					
corrugata, <i>Forbes & Hanl.</i>					
glabra, <i>Forbes & Hanl.</i>					
SALPA, Chamisso.					
runcinata, <i>Cham.</i>					
APPENDICULARIA, Chamisso.					
sp.					

CRUSTACEA.

BRACHYURA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
STENORHYNCHUS, <i>Lamarck</i> .					
phalangium, <i>Pennant</i>					
tenuirostris, <i>Leach</i>					
ACHÆUS, <i>Leach</i> .					
Cranchii, <i>Leach</i>					
INACHUS, <i>Fabr.</i>					
Dorsettensis, <i>Penn.</i>					
dorhynchus, <i>Leach</i>					
leptochirus, <i>Leach</i>					
PISA, <i>Leach</i> (<i>ARCTOPSIS, Lam.</i>).					
tetraodon, <i>Leach</i>					
Gibbsii, <i>Leach</i> (<i>lanata, Lam.</i>) . .					
HYAS, <i>Leach</i> .					
araneus, <i>Fabr.</i>					
coarctatus, <i>Leach</i>					
MAIA, <i>Lam.</i>					
squinado, <i>Herbst</i>					
EURYNOME, <i>Leach</i> .					
aspera, <i>Leach</i>					
XANTHO, <i>Leach</i> .					
florida, <i>Leach</i>					
rivulosa, <i>Edw.</i>					
tuberculata, <i>Couch</i>					
CANCER, <i>Linn.</i>					
pagurus, <i>Linn.</i>					
PILUMNUS, <i>Leach</i> .					
hirtellus, <i>Leach</i>					
PRIMELA, <i>Leach</i> .					
denticulata, <i>Mont.</i>					
CARCINUS, <i>Leach</i> .					
mænas, <i>Linn.</i>					
PORTUMNUS, <i>Leach</i> .					
variegatus, <i>Leach</i> (<i>latipes, Penn.</i>)					
PORTUNUS, <i>Leach</i> .					
puber, <i>Linn.</i>					
corrugatus, <i>Leach</i>					
arcuatus, <i>Leach</i>					
depurator, <i>Leach</i>					
marmoreus, <i>Leach</i>					
holsatus, <i>Fabr.</i>					
pusillus, <i>Leach</i>					
longipes, <i>Risso</i>					
plicatus, <i>Risso</i>					
carcinoides, <i>Kin.</i>					
POLYBIUS, <i>Leach</i> .					
Henslowii, <i>Leach</i>					

CRUSTACEA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
PINNOTHERES, <i>Latr.</i>					
<i>pisum</i> , <i>Penn.</i>					
<i>veterum</i> , <i>Bosc</i>					
GONOPLAX, <i>Leach.</i>					
<i>angulata</i> , <i>Leach</i>					
PLANES, <i>Leach.</i>					
<i>Linnæana</i> , <i>Leach</i>					
EBALIA, <i>Leach.</i>					
<i>Pennantii</i> (<i>tuberosa</i> , <i>Penn.</i>) ..					
<i>Bryerii</i> , <i>Leach</i> (<i>tumefacta</i> , <i>Mont.</i>)					
<i>Cranchii</i> , <i>Leach</i>					
ATELECYCLUS, <i>Leach.</i>					
<i>heterodon</i> , <i>Leach</i> (<i>septemden-</i> <i>tatus</i> , <i>Mont.</i>)					
CORYSTES, <i>Leach.</i>					
<i>Cassivelaunus</i> , <i>Leach</i>					
THIA, <i>Leach.</i>					
<i>polita</i> , <i>Leach</i>					

ANOMOURA.

DROMIA, <i>Edw.</i>					
<i>vulgaris</i> , <i>Edw.</i>					
LITHODES, <i>Latr.</i>					
<i>Maia</i> , <i>Leach</i>					
PAGURUS, <i>Fabr.</i>					
<i>Bernhardus</i> , <i>Linn.</i>					
<i>Prideauxii</i> , <i>Leach</i>					
<i>Cuanensis</i> , <i>Thompson</i>					
<i>Ulidianus</i> , <i>Thompson</i>					
<i>Hyndmanni</i> , <i>Thompson</i>					
<i>lævis</i> , <i>Thompson</i>					
<i>Forbesii</i> , <i>Bell</i>					
<i>Thompsoni</i> , <i>Bell</i>					
<i>fasciatus</i> , <i>Bell</i>					
<i>Dillwynii</i> , <i>Spence Bate</i>					
PORCELLANA, <i>Lamarck.</i>					
<i>platycheles</i> , <i>Penn.</i>					
<i>longicornis</i> , <i>Penn.</i>					
GALATHEA, <i>Fabr.</i>					
<i>squamifera</i> , <i>Leach</i>					
<i>dispersa</i> , <i>Spence Bate</i>					
<i>strigosa</i> , <i>Fabr.</i>					
<i>nexa</i> , <i>Emb.</i>					
<i>Andrewsii</i> , <i>Kinahan</i>					
MUNIDA, <i>Leach.</i>					
<i>Bamfica</i> , <i>Penn.</i> (<i>Rondeletii</i> , <i>Bell</i>)					

CRUSTACEA.

MACROURA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
SCYLLARUS, <i>Fabr.</i>					
<i>arctus</i> , <i>Linn.</i>					
PALINURUS, <i>Fabr.</i>					
<i>Homarus</i> , <i>Linn.</i>					
CALLIANASSA, <i>Leach.</i>					
<i>subterranea</i> , <i>Leach</i>					
GEBIA, <i>Leach.</i>					
<i>stellata</i> , <i>Mont.</i>					
<i>deltura</i> , <i>Leach</i>					
AXIUS, <i>Leach.</i>					
<i>stirhynchus</i> , <i>Leach</i>					
CALOCARIS, <i>Bell.</i>					
<i>Macandreae</i> , <i>Bell</i>					
ASTACUS, <i>Fabr.</i>					
<i>gammarus</i> (<i>L.</i>) (<i>marinus</i> , <i>Fabr.</i>					
<i>vulgaris</i> , <i>Edw.</i>)					
NEPHROPS, <i>Leach.</i>					
<i>Norvegicus</i> , <i>Linn.</i>					
CRANGON, <i>Fabr.</i>					
<i>vulgaris</i> , <i>Fabr.</i>					
<i>fasciatus</i> , <i>Risso</i>					
<i>spinosus</i> , <i>Leach</i>					
<i>sculptus</i> , <i>Bell</i>					
<i>trispinosus</i> , <i>Hailstone</i>					
<i>bispinosus</i> , <i>Westw., Kinahan</i> ..					
<i>Allmanni</i> , <i>Kin.</i>					
<i>Pattersonii</i> , <i>Kin.</i>					
ALPHEUS, <i>Fabr.</i>					
<i>ruber</i> , <i>Edw.</i>					
<i>affinis</i> , <i>Guise</i>					
AUTONOMEA, <i>Risso.</i>					
<i>Olivii</i> , <i>Risso</i>					
NIKA, <i>Risso.</i>					
<i>edulis</i> , <i>Risso</i>					
<i>Couchii</i> , <i>Bell</i>					
ATHANAS, <i>Leach.</i>					
<i>nitescens</i> , <i>Mont., Leach</i>					
HIPPOLYTE, <i>Leach.</i>					
<i>spinus</i> , <i>Sowerby</i>					
<i>varians</i> , <i>Leach</i>					
<i>Craneii</i> , <i>Leach</i>					
<i>Thompsoni</i> , <i>Bell</i>					
<i>Prideauxiana</i> , <i>Leach</i>					
<i>Gordoni</i> , <i>Spence Bate</i>					
<i>fascigera</i> , <i>Gosse</i>					

CRUSTACEA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
HIPPOLYTE, Leach.	fathoms.	fathoms.			
Grayana, Thompson					
Mitchelli, Thompson					
Whitei, Thompson					
Yarrellii, Thompson					
Barleci, Spence Bate					
pandaliformis, Bell					
pusiola, Kröyer					
PANDALUS, Leach.					
Jeffreysii, Spence Bate, Kinahan					
annulicornis, Leach					
leptorhynchus, Kin.					
PALÆMON, Fabr.					
serratus, Penn.					
squilla, Fabr.					
Leachii, Bell					
varians, Leach					
PASIPHEA, Savigny.					
sivado, Risso					
PENÆUS, Fabr.					
caramote, Riess.					

STOMAPODA.

MYSIS, Latr.					
chamæleon, V. Thompson					
vulgaris, V. Thompson					
Griffithsiæ, Bell					
Lamornæ, Couch					
productus, Gosse					
Oberon, Couch					
THYSANOPODA, Edw.					
Couchii, Bell					
MACROMYSIS, White (THEMISTO,					
Goodsir, Bell).					
longispinosus, Goodsir					
brevispinosus, Goodsir					
CYNTHILIA, White (CYNTHIA,					
V. Thomps., Bell).					
Flemingii, Goodsir					
CUMA, Edwards.					
scorpioides, Mont.					
unguiculata, Spence Bate					
VAUNTHOMSONIA, Spence Bate.					
Edwardsii, Kröyer					
crystata, Spence Bate					

CRUSTACEA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
DIASTYLIS, <i>Say</i> (ALAUNA, <i>Goodsir, Bell</i>).					
<i>Rathkii, Kr.</i> (<i>rostrata, Goodsir, Bell</i>)					
EUDORA, <i>Spence Bate</i> .					
<i>truncatula, Spence Bate</i>					
IPHITHOË, <i>Spence Bate</i> (HALIA, <i>Spence Bate, White</i>).					
<i>trispinosa, Goodsir</i>					
BODOTRIA, <i>Goodsir</i> .					
<i>apetosa, Goodsir</i>					
CYRIANASSA, <i>Spence Bate</i> (VENILIA, <i>Spence Bate, White</i>).					
<i>gracilis, Spence Bate</i>					
<i>longicornis, Spence Bate</i>					
SQUILLA, <i>Fabr.</i>					
<i>Desmarestii, Risso</i>					
<i>mantis, Rondelet</i>					
PHYLLOSOMA, <i>Leach</i> .					
<i>Cranchii, Leach</i>					

AMPHIPODA NORMALIA.

TALITRUS, <i>Latr.</i>					
<i>locusta, Auct.</i>					
ORCHESTIA, <i>Leach</i> .					
<i>littorea, Mont.</i>					
<i>Deshayesii, Savig.</i>					
<i>Mediterranea, Costa</i> (<i>lævis, S. Bate</i> ; <i>littorea, var., White</i>)..					
ALLORCHESTES, <i>Dana</i> .					
<i>Nilssonii, Krøyer</i> (<i>Danai, Spence Bate</i>)					
<i>imbricatus, Spence Bate</i>					
NICEA, <i>Nicolet</i> (<i>GALANTHIS, Spence Bate</i>).					
<i>Lubbockiana, Spence Bate</i>					
MONTAGUA, <i>Spence Bate</i> .					
<i>monoculoides, Montagu</i> (<i>Typhis monoculoides, White, Gosse</i>)..					
<i>marina, Spence Bate</i>					
<i>Alderii, Spence Bate</i>					
<i>pollexiana, Spence Bate</i>					
DANAIA, <i>Spence Bate</i> .					
<i>dubia, Spence Bate</i>					
LYSIANASSA, <i>M.-Edw.</i>					
<i>Costæ, M.-Edw.</i>					

CRUSTACEA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
LYSIANASSA, <i>M.-Edw.</i>					
Audouiniana, <i>Spence Bate</i>					
longicornis, <i>Lucas</i> (<i>Chausica</i> , <i>Spence Bate</i> , not <i>M.-Edwards</i>)					
Atlantica, <i>Edw.</i> (<i>marina</i> , <i>Spence</i> <i>Bate</i>)					
CALLISOMA, <i>Hope</i> (<i>SCOPELOCHEIRUS</i> , <i>Spence Bate</i>).					
crenata, <i>Spence Bate</i>					
ANONYX, <i>Kröyer</i> .					
Edwardsii, <i>Kröyer</i>					
minutus, <i>Kröyer</i>					
Holbölli, <i>Kröyer</i>					
ampulla, <i>Kröyer</i>					
denticulatus, <i>Spence Bate</i>					
longipes, <i>Spence Bate</i>					
obesus, <i>Spence Bate</i>					
longicornis, <i>Spence Bate</i>					
OPIS, <i>Kröyer</i> .					
typica, <i>Kröyer</i>					
AMPELISCA, <i>Kröyer</i> (<i>TETROMATUS</i> , <i>Spence Bate</i>).					
Gaimardii, <i>Kröy.</i> (<i>typica</i> , <i>Sp.B.</i>)					
Belliana, <i>Spence Bate</i>					
WESTWOODILLA (<i>WESTWOODIA</i> , <i>Spence Bate</i>).					
cæcula, <i>Spence Bate</i>					
hyalina, <i>Spence Bate</i>					
MONOCULODES, <i>Stimpson</i> .					
carinatus, <i>Spence Bate</i>					
KRÖYERA, <i>Spence Bate</i> .					
arenaria, <i>Spence Bate</i>					
PHOXUS, <i>Kröyer</i> .					
simplex, <i>Spence Bate</i> (<i>Kröyeri</i> , <i>Spence Bate</i> , not <i>Stimpson</i>) . .					
plumosus, <i>Holböll</i>					
Holbölli, <i>Kröy.</i>					
SULCATOR, <i>Spence Bate</i> .					
arenarius, <i>Spence Bate</i>					
UROTHOË, <i>Dana</i> .					
marinus, <i>Spence Bate</i> (<i>Sulcator</i> <i>marinus</i>)					
Bairdii, <i>Spence Bate</i>					
medius, <i>Spence Bate</i>					
elegans, <i>Spence Bate</i>					
GRAYIA, <i>Spence Bate</i> .					
imbricata, <i>Spence Bate</i>					
LILJEBORGIA, <i>Spence Bate</i> .					
pallida, <i>Spence Bate</i>					

CRUSTACEA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
PHLEBRA, <i>Spence Bate</i> .					
<i>antiqua</i> , <i>Spence Bate</i>					
<i>Kinahani</i> , <i>Spence Bate</i>					
ISLEA, <i>M.-Edwards</i> .					
<i>Montagui</i> , <i>M.-Edw.</i>					
IPHIMEDIA, <i>Rathke</i> .					
<i>obesa</i> , <i>Rathke</i>					
<i>Eblanæ</i> , <i>Spence Bate</i>					
OTUS, <i>Spence Bate</i> .					
<i>carinatus</i> , <i>Spence Bate</i>					
ACANTHONOTUS, <i>Owen</i> .					
<i>testudo</i> , <i>Montagu</i>					
DEXAMINE, <i>Leach</i> .					
<i>Loughrimii</i> , <i>Spence Bate</i>					
<i>spinosa</i> , <i>Mont.</i>					
EUSIRUS, <i>Kröyer</i> .					
<i>Edwardi</i> , <i>Spence Bate</i>					
ATYLUS, <i>Leach</i> .					
<i>bispinosus</i> , <i>Spence Bate</i>					
<i>Huxleyanus</i> , <i>Spence Bate</i>					
<i>Gordonianus</i> , <i>Spence Bate</i>					
PHERUSA, <i>Leach</i> .					
<i>cirrus</i> , <i>Spence Bate</i>					
<i>fueicola</i> , <i>Edw.</i>					
CALLIOPE, <i>Leach</i> .					
<i>Leachii</i> , <i>Spence Bate</i>					
EUSIRUS, <i>Spence Bate</i> .					
<i>Helvetiæ</i> , <i>Spence Bate</i>					
LEMBOS, <i>Spence Bate</i> .					
<i>Cambriensis</i> , <i>Spence Bate</i>					
<i>versiculatus</i> , <i>Spence Bate</i>					
<i>Websterii</i> , <i>Spence Bate</i>					
<i>Danmoniensis</i> , <i>Spence Bate</i>					
AÖRA, <i>Kröy.</i> (=LALARIA, <i>Nicolet</i>).					
<i>gracilis</i> , <i>Spence Bate</i>					
EURYSTHEUS, <i>Spence Bate</i> .					
<i>tridentatus</i> , <i>Spence Bate</i>					
<i>tuberculosus</i> , <i>Spence Bate</i>					
GAMMARELLA, <i>Spence Bate</i> .					
<i>brevicaudata</i> , <i>M.-Edw.</i> (=G. or-					
<i>chestiformis</i> , <i>Spence Bate</i>) ..					
CRANGONYX, <i>Spence Bate</i> .					
<i>subterranea</i> , <i>Spence Bate</i>					
AMATHIA, <i>Rathke</i> .					
<i>Sabini</i> , <i>Leach</i>					
GAMMARUS, <i>Fabr.</i>					
<i>locusta</i> , <i>Fabr.</i>					
<i>fluviatilis</i> , <i>Rösel</i>					
<i>gracilis</i> , <i>Rathke</i>					

CRUSTACEA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
GAMMARUS, <i>Fabr.</i>					
<i>camptolops</i> , <i>Leach</i>					
<i>marinus</i> , <i>Leach</i>					
<i>laminatus</i> , <i>Johnston</i>					
<i>longimanus</i> , <i>Leach</i>					
<i>palmatus</i> , <i>Mont.</i> (<i>inæquimanus</i> , <i>Spence Bate</i>)					
<i>grossimanus</i> , <i>Mont.</i>					
<i>maculatus</i> , <i>Johnston</i>					
BATHYPOREIA, <i>Lindström</i> (THER- SITES, <i>Spence Bate</i>).					
<i>pilosa</i> , <i>Lindström</i>					
<i>pelagica</i> , <i>Spence Bate</i>					
<i>Robertsoni</i> , <i>Spence Bate</i>					
LEUCOTHOË, <i>Leach</i> , not <i>Kröyer</i> .					
<i>articulosa</i> , <i>Mont.</i>					
<i>furina</i> , <i>Savig.</i> (<i>procera</i> , <i>Sp. Bate</i>)					
PLEONEXES, <i>Spence Bate</i> .					
<i>gammaroides</i> , <i>Spence Bate</i>					
AMPHITHOË, <i>Leach</i> .					
<i>rubricata</i> , <i>Mont.</i>					
<i>littorina</i> , <i>Spence Bate</i>					
? <i>obtusata</i> , <i>Leach</i>					
? <i>dubia</i> , <i>Johnston</i>					
SUNAMPHITHOË, <i>Spence Bate</i> .					
<i>hamulus</i> , <i>Spence Bate</i>					
<i>conformata</i> , <i>Spence Bate</i>					
PODOCERUS, <i>Leach</i> .					
<i>falcatus</i> , <i>Mont.</i>					
<i>variegatus</i> , <i>Leach</i>					
<i>pulchellus</i> , <i>Leach</i>					
JASSA ?, <i>Leach</i> .					
<i>pelagica</i> , <i>Leach</i>					
SIPHONOCETUS, <i>Kröyer</i> .					
<i>Whitei</i> , <i>Gosse</i>					
ERICHTHONIUS, <i>M.-Edw.</i>					
<i>difformis</i> , <i>M.-Edw.</i>					
CYRTOPHIUM, <i>Dana</i> .					
<i>Darwinii</i> , <i>Spence Bate</i>					
COROPHIUM, <i>Latreille</i> .					
<i>longicornis</i> , <i>Fabr.</i>					
CHELURA, <i>Philippi</i> .					
<i>terebrans</i> , <i>Phil.</i>					
HYPERIA, <i>Latreille</i> .					
<i>Galba</i> , <i>Mont.</i> (<i>Latreillii</i> , <i>Edw.</i> = <i>Metoëcus medusarum</i> , <i>Latr.</i>)					
<i>oblivia</i> , <i>Kröy.</i>					

CRUSTACEA.

AMPHIPODA HYPERINA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
LÆSTRIGONUS, <i>Guérin</i> .					
<i>Fabricii, M.-Edw.</i>					
PHRONIMA, <i>Latr.</i>					
<i>sedentaria, Forsk.</i>					
TYPHIŒ, <i>Risso.</i>					
<i>nolens, Johnston</i>					

AMPHIPODA ABERRANTIA. (LÆMODIPODA of *Latreille*.)

DULICHTIA, <i>Kröyer.</i>					
<i>porrecta, Spence Bate</i>					
<i>falcata, Spence Bate</i>					
PROTO, <i>Leach.</i>					
<i>pedata, Leach</i>					
<i>Goodsirii, Spence Bate</i>					
PROTELLA, <i>Dana.</i>					
<i>longispina, Kröyer</i>					
CAPRELLA, <i>Lamarck.</i>					
<i>linearis, Latr.</i>					
<i>Pennantii, Leach</i>					
<i>tuberculosa, Goodsir</i>					
<i>lobata, Müller</i>					
<i>acuminifera, M.-Edw.</i>					
CYAMUS, <i>Latreille.</i>					
<i>ceti, Linn.</i>					
<i>ovalis, Roussel</i>					
<i>gracilis, Roussel</i>					
<i>Thompsoni, Gosse</i>					

ISOPODA ABERRANTIA. (ANISOPODA of *Dana*.)

ARCTURUS, <i>Latr.</i> (<i>ASTACILLA, Johnston</i> ; <i>LEACHIA, Johnston</i> .)					
<i>longicornis, Sow.</i>					
<i>intermedius, Goodsir</i>					
<i>gracilis, Goodsir</i>					
ANTHURA, <i>Leach.</i>					
<i>gracilis, Mont.</i>					
<i>cylindricus, Mont.</i>					
TANAIS, <i>M.-Edw.</i>					
<i>Dulongii, Aud.</i>					
<i>hirticaudatus, Spence Bate</i>					

CRUSTACEA.

Species.	Range	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
APSEUDES, <i>Leach</i> .					
<i>talpa</i> , <i>Mont.</i>					
ANCEUS, <i>Risso</i> .					
<i>maxillaris</i> , <i>Mont.</i>					
<i>rapax</i> , <i>M.-Edw.</i>					
PRANIZA, <i>Leach</i> .					
<i>ceruleata</i> , <i>Mont.</i>					
<i>fusca</i> ?, <i>Johnston</i>					
<i>Edwardii</i> , <i>Spence Bate</i>					
LIBIOPE, <i>Kröyer</i> .					
<i>balani</i> , <i>Spence Bate</i>					
IONE, <i>Mont.</i>					
<i>thoracica</i> , <i>Mont.</i>					
BOPYRUS, <i>Latr.</i>					
<i>squillarum</i> , <i>Latr.</i>					
<i>hippolytes</i> , <i>Kröyer</i>					
PHRYXUS, <i>Rathke</i> .					
<i>hippolytes</i> , <i>Rathke</i>					
<i>paguri</i> , <i>Rathke</i>					

ISOPODA (NORMALIA).

MUNNA, <i>Kröyer</i> .					
<i>Kröyeri</i> , <i>Goodsir</i>					
<i>Whiteana</i> , <i>Spence Bate</i>					
JERA, <i>Leach</i> .					
<i>albifrons</i> , <i>Leach</i>					
ONISCODA, <i>Latreille</i> .					
<i>maculosa</i> , <i>Leach</i>					
<i>Deshayesii</i> , <i>Lucas</i>					
LIMNORIA, <i>Leach</i> .					
<i>lignorum</i> , <i>Rathke</i> (<i>terebrans</i> , <i>Leach</i>)					
IDOTEA, <i>Fabr.</i>					
<i>pelagica</i> , <i>Leach</i>					
<i>tricuspidata</i> , <i>Desm.</i>					
<i>emarginata</i> , <i>Fabr.</i>					
<i>linearis</i> , <i>Latr.</i>					
<i>acuminata</i> , <i>Leach</i>					
<i>appendiculata</i> , <i>Risso</i>					
LIGIA, <i>Fabr.</i>					
<i>oceanica</i> , <i>Linn.</i>					
SPHÆROMA, <i>Latr.</i>					
<i>serratum</i> , <i>Fabr.</i>					
<i>rugicauda</i> , <i>Leach</i>					
<i>Hookeri</i> , <i>Leach</i>					

CRUSTACEA.

Species.	Range	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
CYMODOCEA, <i>Leach</i> .					
<i>truncata</i> , <i>Leach</i>					
<i>emarginata</i> , <i>Leach</i>					
<i>Montagui</i> , <i>Leach</i>					
<i>rubra</i> , <i>Leach</i>					
<i>viridis</i> , <i>Leach</i>					
NEREA, <i>Leach</i> .					
<i>bidentata</i> , <i>Adams</i>					
CAMPECOPEA, <i>Leach</i> .					
<i>hirsuta</i> , <i>Mont.</i>					
<i>Cranchii</i> , <i>Leach</i>					
CIROLANA, <i>Leach</i> .					
<i>Cranchii</i> , <i>Leach</i>					
EURYDICE, <i>Leach</i> .					
<i>pulchra</i> , <i>Leach</i>					
ÆGA, <i>Leach</i> .					
<i>bicarinata</i> , <i>Leach</i>					
<i>tridens</i> , <i>Leach</i>					
CONILERA, <i>Leach</i> .					
<i>cylindracea</i> , <i>Mont.</i>					
ROCINELA, <i>Leach</i> .					
<i>Danmoniensis</i> , <i>Leach</i>					
<i>monophthalma</i> , <i>Johnston</i>					

ENTOMOSTRACA.

Order I. PHYLLOPODA.

NEBALIA, <i>Leach</i> .					
<i>bipes</i> , <i>O. Fabr.</i>					
ARTEMIA, <i>Leach</i> .					
<i>salina</i> , <i>Linn.</i>					

Order II. CLADOCERA.

EVADNE, <i>Lovén</i> .					
<i>Nordmanni</i> , <i>Lovén</i>					

Order III. OSTRACODA.

Fam. I. Cytheridæ.

CYHERE, <i>Müller</i> .					
<i>flavida</i> , <i>Müll.</i>					
<i>reniformis</i> , <i>Baird</i>					
<i>albo-maculata</i> , <i>Baird</i>					
<i>alba</i> , <i>Baird</i>					

ENTOMOSTRACA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
CY THERE, Müller.					
variabilis, Baird					
aurantia, Baird					
nigrescens, Baird					
Minna, Baird					
angustata, Münster					
acuta, Baird					
pellucida, Baird					
impressa, Baird					
quadridentata, Baird					
convexa, Baird					
CY THEREIS, Rupert Jones.					
Whitei, Baird					
Jonesii, Baird					
antiquata, Baird					
Fam. II. Cypridinidæ.					
CYPRIDINA, M.-Edw.					
Macandrei, Baird					
Brenda, Baird					
Mariæ, Baird					
interpuncta, Baird					
Order IV. COPEPODA.					
Fam. I. Cyclopidæ.					
CANTHOCAMPTUS, Westwood.					
Strömii, Baird					
furcatus, Baird					
minuticornis, Müll.					
ARPACTICUS, M.-Edw.					
chelifer, Müll.					
nobilis, Baird					
ALTEUTHA, Baird.					
depressa, Baird					
Fam. II. Diaptomidæ.					
TEMORA, Baird.					
Finmarchica, Gunner					
CALANUS.					
euchæta, Lubbock					
Anglicus, Lubbock					
ANOMALOCERA, Templeton.					
Patersonii, Temple.					

ENTOMOSTRACA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
Fam. III. Cetochilidæ.					
CETOCHILUS, <i>Vauzème.</i>	fathoms.	fathoms.			
<i>septentrionalis, Goodsir</i>					
PONTELLA, <i>Dana.</i>					
<i>Wollastoni, Lubbock</i>					
PONTELLINA, <i>Dana.</i>					
<i>brevicornis, Lubbock</i>					
PELTIDIUM, <i>Philippi.</i>					
<i>purpureum?, Phil.</i>					
CORYCÆUS, <i>Dana.</i>					
<i>Anglicus, Lubbock</i>					
Fam. IV. Monstrillidæ.					
MONSTRILLA, <i>Dana.</i>					
<i>Anglica, Lubbock</i>					

Species.	On what Animals found.	Observations.
Order V. SIPHONOSTOMA.		
Fam. I. Caligidæ.		
CALIGUS, <i>Müller.</i>		
<i>diaphanus, Nordm.</i>	Various fishes.	
<i>rapax, M.-Edw.</i>	Various fishes.	
<i>Mülleri, Leach</i>	Various fishes.	
<i>centrodonti, Baird</i>	Sea Bream.	
<i>minutus, Otto</i>	Holibut.	
<i>curtus, Müll.</i>	Ray.	
LEPEOPHTHEIRUS, <i>Nordmann.</i>		
<i>Strömii, Baird</i>	On Salmon.	
<i>pectoralis, Müll.</i>	Various fishes.	
<i>Nordmanni, M.-Edw.</i>	Sun-fish.	
<i>hippoglossi, Kröy.</i>	Holibut.	
<i>obscurus, Baird</i>	Brill.	
<i>Thompsoni, Baird</i>	Turbot.	
CHALIMUS, <i>Burmeister.</i>		
<i>scombræ, Burm.</i>	On Mackerel.	
TREBIUS, <i>Kröyer.</i>		
<i>caudatus, Kröy.</i>	On Skate.	
Fam. II. Pandaridæ.		
DINEMOURA, <i>Latreille.</i>		
<i>alata, M.-Edw.</i>	On Shark.	
<i>lamnæ, Johnst</i>	On Shark.	

ENTOMOSTRACA.

Species.	On what Animals found.	Observations.
Fam. III. Cecropidæ.		
PANDARUS, <i>Leach.</i>		
bicolor, <i>Leach</i>	On Shark.	
CECROPS, <i>Leach.</i>		
Latreillei, <i>Leach</i>	On Sun-fish.	
LEMARGUS, <i>Kröyer.</i>		
muricatus, <i>Kröy.</i>	On Sun-fish.	
Fam. IV. Anthosomidæ.		
ANTHOSOMA, <i>Leach.</i>		
Smithii, <i>Leach</i>	On Shark.	
Fam. V. Ergasilidæ.		
NICOTHÖE, <i>M.-Edwards.</i>		
astaci, <i>M.-Edw.</i>	On gills of Lobster	
Fam. VI. Chondracanthidæ.		
CHONDRACANTHUS, <i>De la Roche.</i>		
Zei, <i>De la Roche</i>	Gills of Dory.	
LERNENTOMA, <i>De Blainville.</i>		
cornuta, <i>Müll.</i>	Gills of Sole.	
asellina, <i>L.</i>	Gills of Gurnard.	
lophii, <i>Johnst.</i>	Pouches of Angler.	
LERNEOPODA, <i>De Blainville.</i>		
elongata, <i>Grant</i>	Shark.	
Galei, <i>Kröy.</i>	Shark.	
salmonea, <i>L.</i>	Salmon.	
Fam. VII. Anchorellidæ.		
ANCHORELLA, <i>Cuvier.</i>		
uncinata, <i>Müll.</i>	On the Cod.	
rugosa, <i>Kröy.</i>	On the Cod.	
Fam. VIII. Lerneidæ.		
LERNEA, <i>L.</i>		
branchialis, <i>L.</i>	Gills of Cod.	
LERNEONEMA, <i>M.-Edwards.</i>		
sprattæ, <i>Sby.</i>	On the Sprat.	
Bairdii, <i>Salter</i>	Herring.	
enerasicoli, <i>Turt.</i>	Sprat.	

CIRRIPEDIA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
Fam. I. Balanidæ.					
BALANUS (<i>Lister</i>).					
tintinnabulum, <i>Linn.</i>					
spongiicola, <i>Brown</i>					
perforatus, <i>Bruguère</i>					
Amphitrite, <i>Darwin</i>					
eburneus, <i>Aug. Gould</i>					
improvisus, <i>Darwin</i>					
porcatus, <i>Da Costa</i>					
crenatus, <i>Bruguère</i>					
balanoides, <i>Linn.</i>					
Hameri, <i>Ascanius</i>					
ACASTA, <i>Leach</i> .					
spongites, <i>Poli</i>					
PYRGOMA, <i>Leach</i> .					
Anglicum, <i>G. B. Sby.</i>					
XENOBALANUS, <i>Steenstrup</i> .					
globicipitis, <i>Steenstrup</i>					
CHITHAMALUS, <i>Ranzani</i> .					
stellatus, <i>Poli</i>					
VERRUCA, <i>Schumacher</i> .					
Strömia, <i>O. Müller</i>					
ALCIPPE, <i>Hancock</i> .					
lampas, <i>Hancock</i>					
Fam. II. Lepadidæ.					
LEPAS, <i>Linn.</i>					
anatifera, <i>Linn.</i>					
Hilli, <i>Leach</i>					
anserifera, <i>Linn.</i>					
pectinata, <i>Spengler</i>					
fascicularis, <i>Ellis & Solander</i> . .					
CONCHODERMA, <i>Olfers</i> .					
aurita, <i>Linn.</i>					
virgata, <i>Spengler</i>					
ALEPAS, <i>Sander-Rang</i> .					
parasita, <i>Sander-Rang</i>					
ANELASMA, <i>Darwin</i> .					
squalicola, <i>Lovén</i>					
SCALPELLUM, <i>Leach</i> .					
vulgare, <i>Leach</i>					
POLLICIPES, <i>Leach</i> .					
cornucopia, <i>Leach</i>					

ARACHNIDA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
Order PODOSOMATA.	fathoms.	fathoms.			
Fam. I. Pycnogonidæ.					
PYCNOGONUM, <i>Fabricius</i> . <i>littorale</i> , <i>Strom.</i>					
PHOXICHILUS, <i>Latreille</i> . <i>spinosus</i> , <i>Mont.</i>					
Fam. II. Nymphonidæ.					
PHOXICHILIDIUM, <i>M.-Edwards</i> . <i>coccineum</i> , <i>Johnst.</i> <i>globosum</i> <i>olivaceum</i>					
PALLENE, <i>Johnston</i> . <i>brevirostris</i> , <i>Johnst.</i>					
NYMPHON, <i>Fabricius</i> . <i>gracile</i> , <i>Leach</i> <i>grossipes</i> , <i>O. Fabr.</i> <i>femoratum</i> , <i>Leach</i> <i>pictum</i> <i>giganteum</i> , <i>Johnst.</i>					

ANELLIDA.

** The following list of British Marine Worms is copied, by favour of Dr. J. E. Gray, from an unpublished Catalogue by the late Dr. Johnston.

Order I. TURBELLARIA.					
Fam. I. Planoceridæ.					
LEPTOPLANA, <i>Ehrenberg</i> . <i>subauriculata</i> , <i>Johnston</i> <i>tremellaris</i> , <i>Müller</i> <i>flexilis</i> , <i>Dalyell</i> <i>atomata</i> , <i>Müller</i> <i>ellipsis</i> , <i>Dalyell</i>					
EURYLEPTA, <i>Ehrenberg</i> . <i>cornuta</i> , <i>Müller</i> <i>Dalyellii</i> , <i>Johnston</i> <i>sanguinolenta</i> , <i>Quatrefages</i> <i>vittata</i> , <i>Montagu</i>					
PLANOCERA, <i>Blainville</i> . <i>folium</i> , <i>Grube</i>					

ANELLIDA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
Fam. II. Planariadæ.					
PLANARIA, Müller.					
ulvæ, Oersted					
affinis, Oersted					
alba, Dalyell					
variegata, Dalyell					
? gracilis, Dalyell					
? falcata, Dalyell					
Fam. III. Dalyellidæ.					
TYPHILOPLANA, Ehrenberg.					
flustræ, Dalyell					
CONVOLUTA, Oersted.					
paradoxa, Oersted					
<i>Doubtful species of this family.</i>					
PLANOIDES fusca, Dalyell					}
PLANARIA hirudo, Johnston					
ASTEMMA, Oersted.					
ruffrons, Johnston					
filiformis, Johnston					
CEPHALOTRIX, Oersted.					
lineatus, Dalyell					
flustræ, Dalyell					
TETRASTEMMA, Ehrenberg.					
varicolor, Oersted					
variegatum, Dalyell					
? algæ, Dalyell					
BORLASIA, Johnston.					
olivacea, Johnston					
octoculata, Johnston					
purpurea, Johnston					
Gessensis, Müller					
striata, Rathke					
OMMATOPLEA, Ehrenberg.					
gracilis, Johnston					
rosea, Müller					
alba, Thompson					
melanocephala, Johnston					
pulchra, Johnston					
STYLUS, Johnston.					
viridis, Dalyell					
purpureus, Dalyell					
fragilis, Dalyell					
fasciatus, Dalyell					
LINEUS, T. W. Simmons.					
longissimus, Simmons					

ANNELIDA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
LINEUS, <i>T. W. Simmons</i> .					
<i>gracilis, Goodsir</i>					
<i>lineatus, Johnston</i>					
<i>murenoides, D. Chiaje</i>					
<i>fasciatus, Johnston</i>					
<i>viridis, Dalyell</i>					
<i>albus, Dalyell</i>					
MECKELIA, <i>Leuckart</i> .					
<i>annulata, Montagu</i>					
<i>tænia, Dalyell</i>					
SERPENTARIA, <i>H. D. S. Goodsir</i> .					
<i>fragilis, Goodsir</i>					
<i>fusca, Dalyell</i>					
Order II. BDELLOMORPHA.					
Fam. I. Malacobdellidæ.					
MALACOBDELLA, <i>Blainville</i> .					
<i>grossa, Müller</i>					
<i>Valenciennæi, Blanchard</i>					
<i>anceps, Dalyell</i>					
Order III. BDELLIDEA.					
Fam. I. Branchelliadæ.					
BRANCHELLION, <i>Savigny</i> .					
<i>torpedinis, Savigny</i>					
Fam. II. Piscicolidæ.					
PONTOBDELLA, <i>Leach</i> .					
<i>muricata, Linn.</i>					
<i>verrucata, Grube</i>					
<i>areolata, Leach</i>					
<i>lævis, Blainville</i>					
<i>littoralis, Johnston</i>					
<i>campanulata, Dalyell</i>					
Order IV. SCOLICES.					
Fam. I. Lumbricidæ.					
SENURIS, <i>Hoffmeister</i> .					
<i>lineata, Müller</i>					
CLITELLIO, <i>Savigny</i> .					
<i>arenarius, Müller</i>					
VALLA, <i>Johnston</i> .					
<i>ciliata, Müller</i>					

ANELLIDA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
<p>Order V. GYMNOCOPIA.</p> <p>Fam. I. Tomopteridæ.</p> <p>TOMOPTERIS, <i>Eschscholtz.</i> <i>onisciformis, Grube</i></p>	fathoms.	fathoms.			
<p>Order VI. CHÆTOPODA.</p> <p>Fam. I. Aphroditidæ.</p> <p>APHRODITA, <i>Leach.</i> <i>aculeata, Linn.</i> <i>borealis, Johnston</i> <i>hystrix, Savigny</i></p> <p>LEPIDONOTUS, <i>Leach.</i> <i>squamatus, Linn.</i> <i>clava, Montagu</i> <i>impar, Johnston</i> <i>pharebratus, Johnston</i> <i>cirratus, Fabr.</i> <i>semisculptus, Leach</i> <i>pellucidus, Dyster</i> <i>imbricatus, Linn.</i></p> <p style="text-align: center;"><i>Species not defined.</i></p> <p>APHRODITA <i>squamata, Dalyell</i> . . <i>lepidota, Pallas</i> <i>minuta, Pennant</i> <i>annulata, Pennant</i> <i>velox, Dalyell</i></p> <p>LEPIDONOTUS <i>floccosus?</i>, <i>Dalyell.</i> POLYNOË <i>semisquamosa, Williams</i> }</p> <p>POLYNOË, <i>Oersted.</i> <i>scolopendrina, Savigny</i></p> <p>PHOLOË, <i>Johnston.</i> <i>inornata, Johnston</i> <i>eximia, Dyster</i></p> <p>SIGALION, <i>Aud. & M.-Edwards.</i> <i>boa, Johnston</i></p>					
<p>Fam. II. Amphinomenidæ.</p> <p>EUPHIROSYNE, <i>Savigny.</i> <i>foliosa, Aud. & M.-Edwards.</i> <i>borealis, Oersted</i></p>					
<p>Fam. III. Euniceidæ.</p> <p>EUNICE, <i>Aud. & M.-Edwards.</i> <i>Norvegica, Linn.</i></p>					

ANELLIDA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
<i>EUNICE</i> , <i>Aud. & M.-Edwards</i> .					
<i>annulicornis</i> , <i>Brit. Mus.</i>					
<i>antennata</i> , <i>Savigny</i>					
<i>Harassii</i> , <i>Aud. & M.-Edwards</i>					
<i>sanguinea</i> , <i>Montagu</i>					
<i>margaritacea</i> , <i>Williams</i>					
<i>NORTHIA</i> , <i>Johnston</i> .					
<i>tubicola</i> , <i>Müller</i>					
<i>conchylega</i> , <i>Sars</i>					
<i>LYCIDICE</i> , <i>Savigny</i> .					
<i>Ninetta</i> , <i>Aud. & M.-Edwards</i>					
<i>rufa</i> , <i>Gosse</i>					
<i>LUMBRINERIS</i> , <i>Blainville</i> .					
<i>tricolor</i> , <i>Leach</i>					
Fam. IV. Nereidæ.					
<i>NEREIS</i> , <i>Cuvier</i> .					
<i>brevimana</i> , <i>Johnston</i>					
<i>pelagica</i> , <i>Linn.</i>					
<i>diversicolor</i> , <i>Müller</i>					
<i>cærulea</i> , <i>Linn.</i>					
<i>fimbriata</i> , <i>Müller</i>					
<i>imbecillis</i> , <i>Grube</i>					
<i>Dumerilii</i> , <i>Aud. & M.-Edwards</i>					
<i>pulsatoria</i> , <i>Montagu</i>					
<i>NEREILEPAS</i> , <i>Oersted</i> .					
<i>fucata</i> , <i>Savigny</i>					
<i>HETERONEREIS</i> , <i>Oersted</i> .					
<i>lobulata</i> , <i>Savigny</i>					
<i>renalis</i> , <i>Johnston</i>					
<i>longissima</i> , <i>Johnston</i>					
<i>margaritacea</i> , <i>Johnston</i>					
Fam. V. Nephthyidæ.					
<i>NEPHTHYS</i> , <i>Cuvier</i> .					
<i>cæca</i> , <i>Fabr.</i>					
<i>longisetosa</i> , <i>Oersted</i>					
<i>Homborgii</i> , <i>Cuv.??</i>					
Fam. VI. Phyllodoceidæ.					
<i>PHYLLODOCE</i> , <i>Cuvier</i> .					
<i>lamelligera</i> , <i>Turton</i>					
<i>bilincata</i> , <i>Johnston</i>					
<i>maculata</i> , <i>Linn.</i>					
<i>viridis</i> , <i>Linn.</i>					
<i>ellipsis</i> , <i>Dalyell</i>					

ANELLIDA.

Species.	Range.	Found living at	Ground.	Frequency	Observations.
	fathoms.	fathoms.			
PHYLLODOCE, <i>Cuvier</i> .					
<i>Griffithsii</i> , <i>Johnston</i>					
<i>cordifolia</i> , <i>Dyster</i>					
PSAMATHE, <i>Johnston</i> .					
<i>punctata</i> , <i>Müller</i>					
Fam. VII. Glyceridæ.					
GLYCERA, <i>Savigny</i> .					
<i>mitis</i> , <i>Johnston</i>					
<i>dubia</i> , <i>Blainville</i>					
<i>capitata</i> , <i>Oersted</i>					
<i>nigripes</i> , <i>Johnston</i>					
GONIADA, <i>Aud. & M.-Edwards</i> .					
<i>maculata</i> , <i>Oersted</i>					
Fam. VIII. Syllidæ.					
SYLLIS, <i>Savigny</i> .					
<i>armillaris</i> , <i>Müller</i>					
<i>cornuta</i> , <i>H. Rathke</i>					
<i>prolifera</i> , <i>Müller</i>					
? <i>monoceros</i> , <i>Dalyell</i>					
GATTIOLA, <i>Johnston</i> .					
<i>spectabilis</i> , <i>Johnston</i>					
MYRIANIDA, <i>M.-Edw.</i>					
<i>pinnigera</i> , <i>Montagu</i>					
LOIDA, <i>Johnston</i> .					
<i>macrophthalma</i> , <i>Johnston</i>					
Fam. IX. Amytideidæ.					
AMYTIDEA, <i>Grube</i> .					
<i>maculosa</i> (NEREIS), <i>Montagu</i>					
Fam. X. Ariciadæ.					
NERINE, <i>Johnston</i> .					
<i>vulgaris</i> , <i>Johnston</i>					
<i>coniocephala</i> , <i>Johnston</i>					
<i>Doubtful species.</i> } NERINE contorta (NEREIS), <i>Daly.</i> }					
SPIO, <i>Turton</i> .					
<i>flicornis</i> , <i>Müller</i>					
<i>seticornis</i> , <i>Turton</i>					
<i>crenaticornis</i> , <i>Montagu</i>					
LEUCODORE, <i>Johnston</i> .					
<i>ciliatus</i> , <i>Johnston</i>					

ANELLIDA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
EPHESIA, <i>Rathke</i> .					
<i>gracilis</i> , <i>Rathke</i>					
SPHERODORUM, <i>Oersted</i> .					
<i>peripatus</i> , <i>Johnst</i>					
CIRRATULUS, <i>Lamarck</i> .					
<i>tentaculatus</i> , <i>Mont</i>					
<i>borealis</i> , <i>Lamk</i>					
DODECACERIA, <i>Oersted</i> .					
<i>concharum</i> , <i>Oersted</i>					
Fam. XI. Opheliadæ.					
OPHELIA, <i>Savigny</i> .					
<i>acuminata</i> , <i>Oersted</i>					
AMMOTRYPANE, <i>Rathke</i> .					
<i>limacina</i> , <i>Rathke</i>					
TRAVISIA, <i>Johnston</i> .					
<i>Forbesii</i> , <i>Johnst</i>					
EUMENIA, <i>Oersted</i> .					
<i>crassa</i> , <i>Oersted</i>					
Fam. XII. Siphonostomidæ.					
SIPHONOSTOMA, <i>Cuvier</i> .					
<i>uncinata</i> , <i>Aud. & M.-Edw</i>					
TROPHONIA, <i>Cuvier</i> .					
<i>plumosa</i> , <i>Müller</i>					
Fam. XIII. Telethusidæ.					
ARENICOLA, <i>Savigny</i> .					
<i>piscatorum</i> , <i>Lamk</i>					
<i>branchialis</i> , <i>Aud. & M.-Edw</i> ...					
<i>ecaudata</i> , <i>Johnst</i>					
Fam. XIV. Maldaniadæ.					
CLYMENE, <i>Savigny</i> .					
<i>borealis</i> , <i>Dalyell</i>					
Fam. XV. Terebellidæ.					
TEREBELLA, <i>Montagu</i> .					
<i>conchilega</i> , <i>Pallas</i>					
<i>littoralis</i> , <i>Dalyell</i>					
<i>cirrata</i> , <i>Mont</i>					
<i>nebulosa</i> , <i>Mont</i>					
<i>gigantea</i> , <i>Mont</i>					
<i>constrictor</i> , <i>Mont</i>					
<i>venustula</i> , <i>Mont</i>					
<i>tuberculata</i> , <i>Dalyell</i>					
<i>textrix</i> , <i>Dalyell</i>					
<i>maculata</i> , <i>Dalyell</i>					

ANELLIDA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
VENUSIA, <i>Johnston</i> .					
<i>punctata</i> , <i>Johnst.</i>					
TEREBELLIDES, <i>Sars</i> .					
• <i>Stromii</i> , <i>Sars</i>					
PECTINARIA, <i>Lamarck</i> .					
<i>Belgica</i> , <i>Pallas</i>					
<i>granulata</i> , <i>Linn.</i>					
Fam. XVI. Sabellariadæ.					
SABELLARIA, <i>Lamarck</i> .					
<i>Anglica</i> , <i>Ellis</i>					
<i>crassissima</i> , <i>Lamk.</i>					
<i>lumbricalis</i> , <i>Mont.</i>					
Fam. XVII. Serpulidæ.					
ARIPPASA, <i>Johnston</i> .					
<i>infundibulum</i> , <i>Mont.</i>					
SABELLA, <i>Savigny</i> .					
<i>pavonina</i> , <i>Savigny</i>					
<i>penicillus</i> , <i>Linn.</i>					
<i>vesiculosa</i> , <i>Mont.</i>					
<i>bombyx</i> , <i>Dalyell</i>					
<i>Savignii</i> , <i>Johnst.</i>					
<i>volutacornis</i> , <i>Mont.</i>					
<i>Doubtful Species.</i>					
SABELLA unispira, <i>Sav.</i>					
<i>rosea</i> (AMPHITRITE), <i>Sow.</i> . .					
<i>luna</i> (AMPHITRITE), <i>Dalyell.</i> .					
<i>curta</i> (AMPHITRITE), <i>Mont.</i> . .					
PROTULA, <i>Risso</i> .					
<i>protensa</i> , <i>Philippi</i>					
<i>Dysteri</i> , <i>Huxley</i>					
SERPULA, <i>Linnaeus</i> .					
<i>vermicularis</i> , <i>Ellis</i>					
<i>intricata</i> , <i>Linn.</i>					
<i>reversa</i> , <i>Mont.</i>					
<i>Berkeleyi</i> , <i>Johnst.</i>					
<i>conica</i> , <i>Flem.</i>					
<i>armata</i> , <i>Flem.</i>					
<i>Dysteri</i> , <i>Johnst.</i>					
DITRUPA, <i>Berkeley</i> .					
<i>subulata</i> , <i>Deshayes</i>					
FILOGRANA, <i>Berkeley</i> .					
<i>implexa</i> , <i>Berk.</i>					
OTHOPIA, <i>Johnston</i> .					
<i>Fabricii</i> , <i>Johnst.</i>					

ANELLIDA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
Fam. XVIII. Campontiadae.					
CAMPONTIA, <i>Johnston</i> .					
eruciformis, <i>Johnst.</i>					
Fam. XIX. ? Mæadae.					
MÆA, <i>Johnston</i> .					
mirabilis, <i>Johnst.</i>					
Fam. XX. ? Sipunculidae.					
SYRINX, <i>Bohadsch</i> .					
nudus, <i>Linn.</i>					
papillosus, <i>Thomps.</i>					
Harveii, <i>Forbes</i>					
SIPUNCULUS, <i>Linnaeus</i> .					
Bernhardus, <i>Forbes</i>					
MACRORHYNCHOPTERUS, <i>Rondel.</i>					
Johnstoni, <i>Forbes</i>					
saccatus, <i>Flem.</i>					
tenuicinctus, <i>M. Coy</i>					
Forbesii, <i>M. Coy</i>					
granulosus, <i>M. Coy</i>					
Pallasii, <i>Forbes</i>					
Fam. XXI. Priapulidae.					
PRIAPULUS, <i>Lamarck</i> .					
caudatus, <i>Flem.</i>					
Fam. XXII. Thalassemidæ.					
THALASSEMA, <i>Cuvier</i> .					
Neptuni, <i>Gaertner</i>					
ECHIURUS, <i>Cuvier</i> .					
oxyurus, <i>Pall.</i>					
<i>Species inquirendæ.</i>					
NEREIS, <i>Cuvier</i> .					
iricolor, <i>Mont.</i>					
margarita, <i>Mont.</i>					
lineata, <i>Mont.</i>					
maculosa, <i>Mont.</i>					
rufa, <i>Penn.</i>					
mollis, <i>Linn.</i>					
octentaculata, <i>Mont.</i>					
punctata, <i>Encycl. Méth.</i>					
noctiluca, <i>Linn.</i>					
pinnigera, <i>Mont.</i>					
APHRODITA, <i>Leach</i> .					
annulata, <i>Penn.</i>					
minuta, <i>Penn.</i>					

ANELLIDA.—ENTOZOA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
<i>Species inquirende.</i>					
SPIO, <i>Turton.</i>					
seticornis, <i>Turt.</i>					
crenaticornis, <i>Mont.</i>					
calcareo, <i>Templeton</i>					
BRANCHIARIUS, <i>Mont.</i>					
quadrangularis, <i>Mont.</i>					
DIPLOTIS, <i>Mont.</i>					
hyalina, <i>Mont.</i>					
DERRIS, <i>Adams.</i>					
sanguinea, <i>Adams</i>					

ENTOZOA.

* * * From Dr. Baird's British Museum Catalogue; and Dr. Bellingham's List of Irish Entozoa, in the 'Annals of Natural History,' 1844.

Species.	In what Animals found.	Observations.
Order NEMATOIDEA.		
Fam. Filariadæ.		
FILARIA, <i>Müller.</i>		
? marina, <i>Linn.</i>	Shad and Cod.	
inflexo-caudata, <i>Siebold</i>	Porpoise.	
sp.	Red Gurnard and Mullet.	
TRICHOSOMA, <i>Rudolphi.</i>		
gracilis, <i>Bellingh.</i>	Hake.	
SPIROPTERA, <i>Rudolphi.</i>		
sp.	Skate.	
Fam. Ascaridæ.		
ASCARIS, <i>Linnaeus.</i>		
oseulata, <i>Rud.</i>	Seal.	
aucta, <i>Rud.</i>	Viviparous Blenny.	
rigida, <i>Rud.</i>	Lophius.	
capsularia, <i>Rud.</i>	Cod, &c.	
collaris, <i>Rud.</i>	Flounder, &c.	
clavata, <i>Rud.</i>	Cod, &c.	
constricta, <i>Rud.</i>	Sea-Scorpion, &c.	
rotundata, <i>Rud.</i>	Skate.	

ENTOZOA.

Species.	In what Animals found.	Observations.
<i>ASCARIS, Linnæus.</i>		
<i>acus, Bloch</i>	Herring.	
<i>angulata, Rud.</i>	Lophius.	
<i>tenuissima, Zeder</i>	Whiting.	
<i>succisa, Rud.</i>	Lump-fish.	
Fam. Sclerostomidæ.		
<i>CUCULLANUS, Müller.</i>		
<i>minutus, Rud.</i>	Flounder.	
<i>heterochrous, Rud.</i>	Flounder.	
<i>foveolatus, Lam.</i>	Plaice and Dab.	
<i>STENURUS, Dujardin.</i>		
<i>inflexus, Rud. (part.)</i>	Porpoise.	
<i>PROSTHECOSACTER, Diesing.</i>		
<i>inflexus, Rud.</i>	Porpoise.	
<i>convolutus, Kuhn</i>	Porpoise.	
Order TREMATODA.		
Fam. Onchobothriadæ.		
<i>OCTOBOTHRUM, Leuckart.</i>		
<i>lanceolatum, Leuck.</i>	Shad.	
Fam. Capsalidæ.		
<i>CAPSALA, Bosc.</i>		
<i>coccinea, Cuv.</i>	Sun-fish.	
<i>elongata, Nitzsch</i>	Sturgeon.	
Fam. Distomidæ.		
<i>MONOSTOMA, Zeder.</i>		
<i>flicolle, Rud.</i>	Sea Bream.	
<i>trigonocephalum, Rud.</i>	Turtle.	
<i>DISTOMA, Retzius.</i>		
<i>appendiculatum, Rud.</i>	Shad. &c.	
<i>hispidum, Viborg</i>	Sturgeon.	
<i>megastomum, Rud.</i>	Smooth Shark.	
<i>microcephalum, Baird</i>	Spinous Shark.	
<i>tumidulum, Rud.</i>	Pipe-fish.	
<i>fulvum, Rud.</i>	Skate.	
<i>varicum</i>	Salmon.	
<i>gibbosum?, Rud.</i>	Haddock.	
<i>rufo-viride, Rud.</i>	Conger-eel.	
<i>reflexum?</i>	Cyclopterus.	
<i>excisum, Rud.</i>	Mackerel.	
<i>scabrum, Zeder</i>	Whiting.	
<i>contortum, Rud.</i>	Sun-fish.	
<i>nigro-flavum, Rud.</i>	Sun-fish.	

ENTOZOA.

Species.	In what Animals found.	Observations.
HIRCUDINELLA, <i>Garsin</i> . <i>clavata</i> , <i>Menzies</i>	Bonito.	
Order ACANTHOCEPHALA.		
Fam. Echinorhynchidæ.		
ECHINORHYNCHUS, <i>Müller</i> .		
<i>proteus</i> , <i>Westrumb</i>	Flounder.	
<i>acus</i> , <i>Rud.</i>	Cod, &c.	
<i>gibbosus</i> , <i>Rud.</i>	Herring.	
<i>strumosus</i> , <i>Rud.</i>	Seal.	
Order CESTOIDEA.		
Fam. Rhynchobothridæ.		
RHYNCHOBOTHRUM, <i>Blainville</i> .		
<i>corollatum</i> , <i>Abildg.</i>	Smooth Shark.	
TETRARHYNCHUS, <i>Rudolphi</i> .		
<i>megacephalus</i> , <i>Rud.</i>	Spotted Dog-fish.	
<i>solidus</i> , <i>Drum.</i>	Salmon.	
<i>grossus</i> , <i>Rud.</i>	Salmon.	
<i>rugosus</i> , <i>Baird</i>	Salmon.	
TETRABOTHRIORHYNCHUS, <i>Diesing</i> .		
<i>barbatus</i> , <i>Linn.</i>	Lemon Sole.	
Fam. Tæniadæ.		
BOTHRIOCEPHALUS, <i>Rudolphi</i> .		
<i>fragilis</i> , <i>Rud.</i>	Shad.	
<i>proboscideus</i> , <i>Batsch</i>	Salmon, &c.	
<i>punctatus</i> , <i>Rud.</i>	Turbot, &c.	
<i>tumidulus</i> , <i>Rud.</i>	Ray.	
<i>microcephalus</i> , <i>Rud.</i>	Sun-fish.	
<i>coronatus</i> , <i>Rud.</i>	Skate.	
<i>corollatus</i> , <i>Rud.</i>	Dog-fish.	
<i>paleaceus</i> , <i>Rud.</i>	Dog-fish.	
Fam. Scolecidæ.		
SCOLEX, <i>Müller</i> .		
<i>polymorphus</i> , <i>Rud.</i>	Turbot, &c.	
Order CYSTICA.		
Fam. Cysticidæ.		
ANTHOCEPHALUS, <i>Rudolphi</i> .		
<i>elongatus</i> , <i>Rud.</i>	Sun-fish.	
<i>granulosus</i> ?, <i>Rud.</i>	Whiting, &c.	
<i>paradoxus</i> , <i>Drum.</i>	Turbot.	

ACALEPHIA.

Species.	Where found.	Frequency.	Observations.
Order DISCOPHORA.			
Fam. Sarsiadæ.			
SARSIA, <i>Lesson.</i>			
tubulosa, <i>Sars</i>			
pulchella, <i>Forbes</i>			
gemmifera, <i>Forbes</i>			
prolifera, <i>Forbes</i>			
PLANCIA, <i>Forbes.</i>			
gracilis, <i>Forbes</i>			
BOUGAINVILLIA, <i>Lesson.</i>			
Britannica, <i>Forbes</i>			
nigritella, <i>Forbes</i>			
pyramidata, <i>Gosse?</i>			
crucifera, <i>Gosse?</i>			
simplex, <i>Gosse?</i>			
LIZZIA, <i>Forbes.</i>			
octopunctata, <i>Forbes</i>			
blondina, <i>Forbes</i>			
MODEERIA, <i>Forbes.</i>			
formosa, <i>Forbes</i>			
EUPHYSA, <i>Forbes.</i>			
aurata, <i>Forbes</i>			
STEENSTRUPIA, <i>Forbes.</i>			
rubra, <i>Forbes</i>			
flavcola, <i>Forbes</i>			
Fam. Geryoniadæ.			
GERYONIA, <i>Péron.</i>			
appendiculata, <i>Forbes</i>			
TIMA, <i>Eschscholtz.</i>			
Bairdii, <i>Johnst.</i>			
GERYONOPSIS, <i>Forbes.</i>			
delicatula, <i>Forbes</i>			
THAUMANTIAS, <i>Eschscholtz.</i>			
pilosella, <i>Forbes</i>			
quadrata, <i>Forbes</i>			
aëronautica, <i>Forbes</i>			
octona, <i>Forbes</i>			
maculata, <i>Forbes</i>			
melanops, <i>Forbes</i>			
globosa, <i>Forbes</i>			
convexa, <i>Forbes</i>			
gibbosa, <i>Forbes</i>			

ACALEPHA.

Species.	Where found.	Frequency.	Observations.
<i>THAUMANTIAS, Eschscholtz.</i>			
<i>lineata, Forbes</i>			
<i>pileata, Forbes</i>			
<i>Sarnica, Forbes</i>			
<i>Thompsoni, Forbes</i>			
<i>hemisphærica, Linn.</i>			
<i>inconspicua, Forbes</i>			
<i>punctata, Forbes</i>			
<i>lucifera, Forbes</i>			
<i>Buskiana, Gosse?</i>			
<i>corynetes, Gosse?</i>			
<i>undulata, Gosse?</i>			
<i>confluens, Gosse?</i>			
<i>SLABBERIA, Forbes.</i>			
<i>halterata, Forbes</i>			
<i>catenata, Gosse?</i>			
Fam. Oceaniadæ.			
<i>TURRIS, Less.</i>			
<i>digitalis, Fabr.</i>			
<i>neglecta, Less.</i>			
<i>SAPHENIA, Eschscholtz.</i>			
<i>dinema, Esch.</i>			
<i>Titania, Gosse?</i>			
<i>OCEANIA, Péron.</i>			
<i>octona, Flem.</i>			
<i>episcopalis, Forbes</i>			
<i>ducalis, Gosse?</i>			
<i>turrita, Forbes</i>			
<i>globulosa, Forbes</i>			
<i>pusilla, Gosse?</i>			
Fam. Circeadæ.			
<i>CIRCE, Mertens.</i>			
<i>rosea, Forbes</i>			
Fam. Æquoreadæ.			
<i>STOMOBRACTIUM, Brandt.</i>			
<i>octocostatum, Less.</i>			
<i>POLYXENIA, Eschscholtz.</i>			
<i>Alderi, Forbes</i>			
<i>ÆQUOREA, Péron.</i>			
<i>Forskallii, Gosse?</i>			
<i>vitrina, Gosse?</i>			
<i>Forbesiana, Gosse?</i>			

ACALEPHA.

Species.	Where found.	Frequency.	Observations.
Fam. Willsiadæ.			
WILLSIA, <i>Forbes</i> . stellata, <i>Forbes</i>			
Fam. Medusadæ.			
MEDUSA, <i>Linn.</i> aurita, <i>Linn.</i> campanula, <i>Lamk.</i> surireæ, <i>Lamk.</i>			
CYANÆA, <i>Péron.</i> capillata, <i>Linn.</i> Lamarekii, <i>Dicq.</i>			
CHRYSAORA, <i>Péron.</i> hyoscella, <i>Linn.</i> cyclonota, <i>Péron</i>			
PELAGIA, <i>Péron.</i> cyanella, <i>Lamk.</i>			
Fam. Rhizostomadæ.			
CASSIOPEA, <i>Péron.</i> lunulata, <i>Esch.</i>			
RHIZOSTOMA, <i>Cuvier.</i> pulsus, <i>Gmel.</i>			
Order CTENOPHORA.			
Fam. Callianiradæ.			
CYDIPPE, <i>Esch.</i> pileus, <i>Müll.</i> Flemingii, <i>Forbes</i> infundibulum, <i>Müll.</i> lagena, <i>Forbes</i> pomiformis, <i>Paterson</i>			
BOLINA, <i>Paterson.</i> Hibernica, <i>Paterson</i>			
Fam. Beroidæ.			
BEROË, <i>Müller.</i> cucumis, <i>Fabr.</i> fulgens, <i>Flem.</i> borealis, <i>Less.</i>			
ALCINOË, <i>Cuv.</i> rotunda Smithii			

ACALEPHIA.—ECHINODERMATA.

Species.	Where found.	Frequency.	Observations.
Order SIPHONOPHORA.			
Fam. Diphyidæ.			
DIPHYES, <i>Cuvier.</i>			
<i>campanulifera, Esch.</i>			
Fam. Physaliadæ.			
PHYSALIA, <i>Lamarck.</i>			
<i>pelagica, Esch.</i>			
Fam. Velelladæ.			
VELELLA, <i>Lamarck.</i>			
<i>vulgaris, Leach.</i>			
RAFARIA, <i>Esch.</i>			
<i>pacillum, Mont.</i>			

CLASS ECHINODERMATA.

SUBCLASS HYPOSTOMATA.

Species.	Range	Found living at	Ground.	Frequency.	Observations.
Order CRINOIDEA.					
COMATULA, <i>Lamarck.</i>					
<i>rosacea, Link</i>	fathoms.	fathoms.			
<i>Celtica, Barrett</i>					
Order OPHIUROIDEA.					
Fam. Ophiuriadæ.					
OPHIURA, <i>Lamarck.</i>					
<i>texturata, Lamk.</i>					
<i>albida, Forbes</i>					
OPHIOCOMA, <i>Agassiz.</i>					
<i>neglecta, Johnst.</i>					
<i>punctata, Forbes</i>					
<i>filiformis, Müller</i>					
<i>securigera, D. & K.</i>					
<i>bellis, Link</i>					
<i>brachiata, Mont.</i>					
<i>Ballii, Thomps.</i>					
<i>Goodsiri, Forbes</i>					
<i>granulata, Link</i>					
<i>rosula, Link</i>					
<i>minuta, Forbes.</i>					

ECHINODERMATA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
Subfam. Euryalina.					
ASTROPHYTON, <i>Link.</i>					
scutatum, <i>Link</i>					
Order ASTEROIDEA.					
Fam. Asteriadae.					
URASTER, <i>Agassiz.</i>					
glacialis, <i>Linn.</i>					
rubens, <i>Linn.</i>					
violacea, <i>Müller</i>					
hispida, <i>Penn.</i>					
rosca, <i>Müller</i>					
ECHINASTER, <i>Müller & Troschel.</i>					
oculatus, <i>Penn.</i>					
SOLASTER, <i>Forbes.</i>					
endeca, <i>Linn.</i>					
papposa, <i>Linn.</i>					
PALMIPES, <i>Link.</i>					
membranaceus, <i>Retz.</i>					
ASTERINA, <i>Nardo.</i>					
gibbosa, <i>Penn.</i>					
GONIASTER, <i>Agassiz.</i>					
Templetoni, <i>Thomps.</i>					
equestris, <i>Gmelin</i>					
ASTERIAS, <i>Linnaeus.</i>					
aurantiaca, <i>Linn.</i>					
LUIDIA, <i>Forbes.</i>					
fragilissima, <i>Forbes</i>					
Savignii, <i>Audouin</i>					
Order ECHINOIDEA.					
CIDARIS, <i>Leske.</i>					
papillata, <i>Leske</i>					
ECHINUS, <i>Linnaeus.</i>					
sphaera, <i>Müller</i>					
Flemingii, <i>Ball</i>					
miliaris, <i>Leske</i>					
lividus, <i>Lamk.</i>					
melio, <i>Lamk.</i>					
Norvegicus, <i>D. & K.</i>					
neglectus, <i>Lamk.</i>					
Order CLYPEASTEROIDEA.					
ECHINOCYAMUS, <i>Leske.</i>					
pusillus, <i>Müller</i>					

ECHINODERMATA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
ECHINARACHNIUS, <i>Leske</i> . placenta, <i>Gmelin</i>					
Order SPATANGOIDEA.					
SPATANGUS, <i>Klein</i> . purpureus, <i>Müller</i>					
BRISSUS, <i>Klein</i> . lyrifer, <i>Forbes</i>					
AMPHIDOTUS, <i>Agassiz</i> . cordatus, <i>Penn</i> roseus, <i>Forbes</i> gibbosus, <i>Barrett</i>					
Order HOLOTHUROIDEA.					
Fam. Pentactidæ.					
CUCUMARIA, <i>Cuvier</i> . frondosa, <i>Gunner</i> ? fucicola, <i>Forbes & Goodsir</i> pentactes, <i>Müller</i> ? Montagui, <i>Flem.</i> ? Neilli, <i>Flem.</i> ? dissimilis, <i>Flem.</i> fusiformis, <i>Forbes & Goodsir</i> Hyndmanni, <i>Thomps.</i>					
OENUS, <i>Forbes</i> (= CUCUMARIA?). brunneus, <i>Forbes</i> lacteus, <i>Forbes & Goodsir</i>					
PSOLINUS, <i>Forbes</i> . brevis, <i>Forbes & Goodsir</i>					
Fam. Thyonidæ.					
THYONE, <i>Oken</i> . fusus, <i>Müll.</i> (papillosa, <i>Abildg.</i>) raphanus, <i>Duben & Koren</i> communis, <i>F. & G.</i> (THYONIDIUM, <i>D. & K.</i>) Portlockii, <i>Forbes</i> Drummondii, <i>Thomps.</i> pellucidum, <i>Fahl.</i> (<i>C. hyalina, F.</i>)					
HOLOTHURIA, <i>Linn.</i> nigra, <i>Couch</i> intestinalis tubulosa, <i>Linn.</i>					

ECHINODERMATA.—POLYZOA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
Fam. Psolidæ.					
<i>PSOLUS</i> , <i>Oken</i> .					
phantopus, <i>Linn.</i>					
Forbesii					
Fam. Synaptidæ.					
<i>SYNAPTA</i> , <i>Esch.</i>					
inhærens, <i>Müll.</i>					
digitata, <i>Mont.</i>					

POLYZOA.

. From the British Museum Catalogue by Mr. George Busk.

Order I. INFUNDIBULATA.					
Suborder I. CHEILOSTOMATA.					
Fam. II. Salicornariadæ.					
<i>SALICORNARIA</i> , <i>Cuvier</i> .					
farciminoïdes, <i>Johnst.</i>					
Fam. III. Cellulariadæ.					
<i>CELLULARIA</i> , <i>Pallas</i> .					
Peachii, <i>Busk</i>					
<i>MENIPEA</i> , <i>Lamouroux</i> .					
ternata, <i>Ellis & Soland.</i>					
<i>SCRUPOCELLARIA</i> , <i>Van Beneden</i> .					
scrupæa, <i>Busk</i>					
scruposa, <i>Linn.</i>					
<i>CANDA</i> , <i>Lamouroux</i> .					
reptans, <i>Pall.</i>					
Fam. IV. Scrupariadæ.					
<i>SCRUPARIA</i> , <i>Oken</i> .					
chelata, <i>Linn.</i>					
<i>SALPINGIA</i> , <i>Coppin</i> .					
Hassallii, <i>Coppin</i>					
<i>HIPPOTHOA</i> , <i>Lamouroux</i> .					
catenularia, <i>Jameson</i>					
divaricata, <i>Lamx.</i>					

POLYZOA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
<i>ÆTEA, Lamourouæ.</i>					
<i>anguina, Linn.</i>					
<i>truncata, Landsb.</i>					
<i>BEANIA, Johnston.</i>					
<i>mirabilis, Johnst.</i>					
Fam. VI. Gemellariadæ.					
<i>GEMELLARIA, Savigny.</i>					
<i>loricata, Linn.</i>					
<i>NOTAMIA, Fleming.</i>					
<i>bursaria, Linn.</i>					
Fam. VII. Cabereadæ.					
<i>CABEREA, Lamourouæ.</i>					
<i>Hookeri, Flem.</i>					
Fam. VIII. Bicellariadæ.					
<i>BICELLARIA, De Blainville.</i>					
<i>ciliata, Linn.</i>					
<i>BUGULA, Oken.</i>					
<i>neritina, Linn.</i>					
<i>flabellata, J. V. Thomps.</i>					
<i>avicularia, Pall.</i>					
<i>plumosa, Pall.</i>					
<i>Murrayana, Bean.</i>					
Fam. IX. Flustradæ.					
<i>FLUSTRA, Linn.</i>					
<i>foliacea, Linn.</i>					
<i>papyracea, Ellis</i>					
<i>truncata, Linn.</i>					
<i>CARBASEA, Gray.</i>					
<i>papyrea, Pall.</i>					
Fam. X. Membraniporidæ.					
<i>MEMBRANIPORA, De Blainville.</i>					
<i>membranacea, Linn.</i>					
<i>pilosa, Pall.</i>					
<i>coriacea, Esper</i>					
<i>lineata, Linn.</i>					
<i>Flemingii, Busk</i>					
<i>Rossellii, Audouin</i>					
<i>Laeroixii, Savigny</i>					
<i>monostachys, Busk</i>					

POLYZOA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
<i>LEPRALIA, Johnston.</i>					
<i>Brongniartii, Aud.</i>					
<i>Landsborovii, Johnst.</i>					
<i>reticulata, Macgillivray</i>					
<i>auriculata, Hassall</i>					
<i>concinna, Busk</i>					
<i>verrucosa, Esper</i>					
<i>violacea, Johnst.</i>					
<i>spinifera, Johnst.</i>					
<i>trispinosa, Johnst.</i>					
<i>coccinea, Abildg.</i>					
<i>linearis, Hassall</i>					
<i>eliata, Pall.</i>					
<i>Gattyæ, Landsb.</i>					
<i>Hyndmanni, Johnst.</i>					
<i>variolosa, Johnst.</i>					
<i>nitida, Fabr.</i>					
<i>annulata, Fabr.</i>					
<i>bispinosa, Johnst.</i>					
<i>Peachii, Johnst.</i>					
<i>ventricosa, Hassall</i>					
<i>melolontha, Landsb.</i>					
<i>innominata, Couch</i>					
<i>punctata, Hassall</i>					
<i>figularis, Johnst.</i>					
<i>pertusa, Esper</i>					
<i>Pallasiana</i>					
<i>labrosa, Busk</i>					
<i>simplex, Johnst.</i>					
<i>Malusii, Aud.</i>					
<i>granifera, Johnst.</i>					
<i>hyalina, Linn.</i>					
Fam. XI. <i>Celleporidæ.</i>					
<i>CELLEPORA, Fabr.</i>					
<i>pumicosa, Linn.</i>					
<i>Hassallii, Johnst.</i>					
<i>vitrina, Couch</i>					
<i>ramulosa, Linn.</i>					
<i>Skenei, Ellis & Soland.</i>					
Fam. XII. <i>Escharidæ.</i>					
<i>ESCHARA, Ray.</i>					
<i>foliacea, Ellis & Soland.</i>					
<i>cervicornis, Soland.</i>					
<i>RETEPORA, Lamarck.</i>					
<i>cellulosa, Linn.</i>					
<i>Beaniana, King</i>					

POLYZOA.

Species.	Range	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
Suborder II. CYCLOSTOMATA.					
Fam. I. Tubuliporidae.					
TUBULIPORA, <i>Lamarck</i> .					
patina, <i>Linn.</i>					
hispida, <i>Flem.</i>					
penicillata, <i>Johnst.</i>					
truncata, <i>Jameson</i>					
lobulata, <i>Hassall</i>					
phalangea, <i>Couch</i>					
flabellaris, <i>Fabr.</i>					
serpeus, <i>Linn.</i>					
hyalina, <i>Couch</i>					
DIASTOPORA, <i>Lamouroux</i> .					
obelia, <i>Flem.</i>					
IDMONEA, <i>Lamouroux</i> .					
Atlantica, <i>Forbes</i>					
PUSTULIPORA, <i>De Blainville</i> .					
proboscidea, <i>M.-Edw.</i>					
deflexa, <i>Couch</i>					
ALECTO, <i>Lamouroux</i> .					
granulata, <i>M.-Edw.</i>					
major, <i>Johnst.</i>					
dilatans, <i>Johnst.</i>					
Fam. II. Crisiadae.					
CRISIA, <i>Lamouroux</i> .					
eburnea, <i>Linn.</i>					
denticulata, <i>Lamk.</i>					
aculeata, <i>Hassall</i>					
geniculata, <i>M.-Edw.</i>					
CRISIDIA, <i>M.-Edwards</i> .					
cornuta, <i>Linn.</i>					
setacea, <i>Couch</i>					
Suborder III. CTENOSTOMATA.					
Fam. I. Alcyonidiadae.					
ALCYONIDIUM, <i>Lamouroux</i> .					
gelatinosum, <i>Pallus</i>					
hirsutum, <i>Flem.</i>					
parasiticum, <i>Flem.</i>					
Cycloum, <i>Hass.</i>					
papillosum, <i>Hass.</i>					
SARCOCHITUM, <i>Hass.</i>					
polyoum, <i>Johnst.</i>					

POLYZOA.—ZOOPHYTA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
Fam. II. Vesiculariadae.					
AMATHIA, <i>Lamouroux.</i>	fathoms.	fathoms.			
<i>lendigera</i> , <i>Linn.</i>					
VESICULARIA, <i>Thompson.</i>					
<i>spinosa</i> , <i>Linn.</i>					
VALKERIA, <i>Fleming.</i>					
<i>cuscuta</i> , <i>Ellis</i>					
<i>uva</i> , <i>Linn.</i>					
<i>pustulosa</i> , <i>Johnst.</i>					
MIMOSELLA, <i>Hincks.</i>					
<i>gracilis</i>					
AVENELLA, <i>Dalyell.</i>					
<i>Dalyellii</i>					
NOTELLA, <i>Gosse.</i>					
<i>stipata</i> , <i>Gosse</i>					
BOWERBANKIA, <i>Farre.</i>					
<i>imbricata</i> , <i>Johnst.</i>					
FARRELLA, <i>Ehrenberg.</i>					
<i>repens</i> , <i>Johnst.</i>					
<i>elongata</i>					
<i>gigantea</i>					
ANGUINELLA, <i>Van Beneden.</i>					
<i>palmata</i>					
Fam. III. Pedicellinidae.					
PEDICELLINA, <i>Sars.</i>					
<i>echinata</i> , <i>Sars</i>					
<i>Belgica</i>					
<i>gracilis</i>					

ZOOPHYTA.

. This list is taken from Johnston's "British Zoophytes," and Gosse's "Actinologia."

Order HYDROIDA.					
Fam. I. Corynidae.					
CLAVA, <i>Gmelin.</i>					
<i>multicornis</i> , <i>Pall.</i>					
HYDRACTINIA, <i>Van Beneden.</i>					
<i>echinata</i> , <i>Flem.</i>					
<i>Alderi</i> , <i>Johnst.</i>					
<i>littoralis</i> , <i>Johnst.</i>					
MYRIOTHELA, <i>Sars.</i>					
<i>arctica</i>					

ZOOPHYTA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
<i>CORYNE</i> , <i>Gaertner</i> .					
<i>pusilla</i> , <i>Pall</i>					
<i>ramosa</i> , <i>Fubr.</i>					
<i>sessilis</i>					
<i>stauridia</i>					
<i>Cerberus</i>					
Fam. II. Tubulariadae.					
<i>EUDENDRIUM</i> , <i>Ehrenberg</i> .					
<i>ramosum</i> , <i>Ellis</i>					
<i>rameum</i> , <i>Pall.</i>					
<i>TUBULARIA</i> , <i>Linnaeus</i> .					
<i>indivisa</i> , <i>Linn.</i>					
<i>Dumortierii</i> , <i>Van Beneden</i>					
<i>larynx</i> , <i>Ellis</i>					
<i>gracilis</i> , <i>Harvey</i>					
<i>polyceps</i> , <i>Dalyell</i>					
<i>CORYMORPHA</i> , <i>Sars</i> .					
<i>nutans</i> , <i>Sars</i>					
Fam. III. Sertulariadae.					
<i>HALECIUM</i> , <i>Oken</i> .					
<i>halecinum</i> , <i>Ellis</i>					
<i>Beanii</i> , <i>Johnst.</i>					
<i>muricatum</i> , <i>Ellis & Soland.</i>					
<i>SERTULARIA</i> , <i>Linnaeus</i> .					
<i>polyzonias</i> , <i>Linn.</i>					
<i>rugosa</i> , <i>Ellis</i>					
<i>rosacea</i> , <i>Linn.</i>					
<i>pumila</i> , <i>Ray</i>					
<i>Evansii</i> , <i>Ellis & Soland.</i>					
<i>nigra</i> , <i>Jameson</i>					
<i>pinnata</i> , <i>Pall.</i>					
<i>pinaster</i> , <i>Ellis & Soland.</i>					
<i>Margareta</i> , <i>Hassall</i>					
<i>fallax</i> , <i>Flem.</i>					
<i>tamarisea</i> , <i>Ellis</i>					
<i>abietina</i> , <i>Linn.</i>					
<i>filicula</i> , <i>Ellis & Soland.</i>					
<i>operculata</i> , <i>Linn.</i>					
<i>argentea</i> , <i>Ellis & Soland.</i>					
<i>cupressina</i> , <i>Ellis</i>					
<i>gracilis</i>					
<i>RETICULARIA</i> , <i>W. Thompson</i> .					
<i>immersa</i>					
<i>COPPINIA</i> , <i>Hassall</i> .					
<i>arcta</i>					

ZOOPHYTA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
THUIARIA, Fleming.					
<i>thuia, Linn.</i>					
<i>articulata, Flem.</i>					
ANTENNULARIA, Lamarck.					
<i>antennina, Linn.</i>					
<i>ramosa, Lamæ.</i>					
PLUMULARIA, Lamarck.					
<i>falcata, Linn.</i>					
<i>cristata, Lamk.</i>					
<i>pennatula, Ellis & Soland.</i>					
<i>pinnata, Lamk.</i>					
<i>setacea, Ellis</i>					
<i>Catharina, Johnst.</i>					
<i>echinulata, Lamk.</i>					
<i>fascis, Dalyell</i>					
<i>myriophyllum, Linn.</i>					
<i>frutescens, Ellis & Soland.</i>					
Fam. IV. Campanulariadae.					
LAOMEDEA, Lamouroux.					
<i>dichotoma, Linn.</i>					
<i>geniculata, Linn.</i>					
<i>gelatinosa, Pall.</i>					
<i>obliqua, Saunders</i>					
<i>Flemingii, M.-Edw.</i>					
<i>lacerata</i>					
CAMPANULARIA, Lamarck.					
<i>volubilis, Linn.</i>					
<i>integra, Macgilliv.</i>					
<i>lacerata, Johnst.</i>					
<i>lævis, Couch.</i>					
<i>serpens</i>					
<i>intertexta, Couch</i>					
<i>syringa, Linn.</i>					
<i>parvula</i>					
<i>caliculata</i>					
<i>verticillata, Linn.</i>					
<i>dumosa, Flem.</i>					
Order ACTINOIDA.					
Fam. Actiniadae.					
ACTINOLOBIA, Blainville.					
<i>dianthus, Blainv.</i>					
SAGARTIA, Gosse.					
<i>bellis, Ellis</i>					
<i>miniata, Gosse</i>					
<i>rosca, Gosse</i>					

ZOOPHYTA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
SAGARTIA, <i>Gosse.</i>					
<i>ornata, Holdsworth.</i>					
<i>ichthystoma, Gosse</i>					
<i>nivca, Gosse</i>					
<i>sphyrodeta, Gosse</i>					
<i>pallida, Holdsworth</i>					
<i>pura, Alder</i>					
<i>coccinea, Müll.</i>					
<i>troglydites, Johnst.</i>					
<i>viduata, Müll.</i>					
<i>parasitica, Johnst.</i>					
<i>chrysosplenium, Cocks.</i>					
ADAMSIA, <i>Forbes.</i>					
<i>palliata, Forbes.</i>					
PHELLIA, <i>Gosse.</i>					
<i>murocincta, Gosse.</i>					
<i>gausapata, Gosse</i>					
<i>pieta, Gosse</i>					
<i>Brodricii, Gosse</i>					
GREGORIA, <i>Gosse.</i>					
<i>fenestrata, Gosse</i>					
AIPTASIA, <i>Cocks.</i>					
<i>Couchii, Cocks</i>					
ANTHEA, <i>Gaërtner.</i>					
<i>cercus, Ellis</i>					
ACTINIA, <i>Linn.</i>					
<i>mesembryanthemum, Ellis.</i>					
BOLOCERA, <i>Johnst.</i>					
<i>Tuediæ, Johnst.</i>					
<i>eques, Gosse</i>					
BUNODES, <i>Gosse.</i>					
<i>gemmacca, Ellis</i>					
<i>thallia, Gosse</i>					
<i>Ballii, Cocks.</i>					
<i>coronata, Gosse.</i>					
TEALIA, <i>Gosse.</i>					
<i>digitata, Müll.</i>					
<i>tuberculata, Cocks</i>					
<i>crassicornis, Müll.</i>					
HORMATHIA, <i>Gosse.</i>					
<i>margarita, Gosse</i>					
STOMPHIA, <i>Gosse.</i>					
<i>Churchiæ, Gosse</i>					
ILYANTHUS, <i>Forbes.</i>					
<i>Scoticus, Forbes</i>					
<i>Mitchellii, Gosse</i>					
PEACHIA, <i>Gosse.</i>					
<i>hastata, Gosse</i>					
<i>undata, Gosse</i>					

ZOOPHYTA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
PEACHIA, <i>Gosse</i> .					
<i>triphylla</i> , <i>Gosse</i>					
<i>cylindrica</i> , <i>Reid</i>					
HALCAMP, <i>Gosse</i> .					
<i>chrysanthemum</i> , <i>Peach</i>					
<i>microps</i> , <i>Gosse</i>					
EDWARDSIA, <i>Quatrefages</i> .					
<i>callimorpha</i> , <i>Gosse</i>					
<i>carnea</i> , <i>Gosse</i>					
<i>Beautempsii</i> , <i>Quatref.</i>					
ARACHNACTIS, <i>Sars</i> .					
<i>albida</i> , <i>Sars</i>					
CERIANTHUS, <i>J. Haime</i> .					
<i>Lloydii</i> , <i>Gosse</i>					
<i>vermicularis</i> , <i>Forbes</i>					
CAPNEA, <i>Forbes</i> .					
<i>sanguinea</i> , <i>Forbes</i>					
AURELIANIA, <i>Gosse</i> .					
<i>angusta</i> , <i>Gosse</i>					
<i>heterocera</i> , <i>Thomps.</i>					
CORYNACTIS, <i>Allman</i> .					
<i>viridis</i> , <i>Allman</i>					
Fam. Zoanthidæ.					
ZOANTHUS, <i>Cuvier</i> .					
<i>Couchii</i> , <i>Johnst.</i>					
<i>sulcatus</i> , <i>Gosse</i>					
<i>Alderi</i> , <i>Gosse</i>					
Fam. Caryophylleadæ.					
CYATHINA, <i>Ehrenberg</i> .					
<i>Smithii</i> , <i>Flem.</i>					
PARACYATHUS, <i>M.-Edw.</i>					
<i>Taxilianus</i> , <i>Gosse</i>					
<i>Thulensis</i> , <i>Gosse</i>					
<i>pteropus</i> , <i>Gosse</i>					
DESMOPHYLLUM, <i>Ehrenberg</i> .					
<i>Stokesii</i> , <i>M.-Edw.</i>					
SPIENOTROCHUS, <i>M.-Edwards</i> .					
<i>Macandrewanus</i> , <i>M.-Edw.</i>					
<i>Wrightii</i> , <i>Gosse</i>					
ULOCYATHUS, <i>Sars</i> .					
<i>arcticus</i> , <i>Sars</i>					
OCULINA, <i>Lamarck</i> .					
<i>prolifera</i> , <i>Linn.</i>					
HOPLANGIA, <i>Gosse</i> .					
<i>Durotrix</i> , <i>Gosse</i>					

ZOOPHYTA.—FORAMINIFERA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
BALANOPHYLLIA, <i>Wood</i> .					
<i>regia</i> , <i>Gosse</i>					
Suborder ALCYONARIA.					
Fam. I. Pennatuladæ.					
PENNATULA, <i>Linnaeus</i> .					
<i>phosphorea</i> , <i>Linn.</i>					
VIRGULARIA, <i>Lamarck</i> .					
<i>mirabilis</i> , <i>Linn.</i>					
PAVONARIA, <i>Cuvier</i> .					
<i>quadrangularis</i> , <i>Pall.</i>					
Fam. II. Alcyonidæ.					
ALCYONIUM, <i>Linnaeus</i> .					
<i>digitatum</i> , <i>Linn.</i>					
<i>glomeratum</i>					
SARCODICTYON, <i>Forbes</i> .					
<i>catenata</i> , <i>Forbes</i>					
<i>agglomerata</i>					
Fam. III. Gorgoniadæ.					
GORGONIA, <i>Linnaeus</i> .					
<i>verrucosa</i> , <i>Linn.</i>					
<i>pinnata</i> , <i>Linn.</i>					
<i>anceps</i> , <i>Ellis</i>					
PRIMNOA, <i>Lamarck</i> .					
<i>lepadifera</i> , <i>Linn.</i>					

FORAMINIFERA.

. This list of British Foraminifera is taken from Prof. Williamson's "Recent Foraminifera of Great Britain," published by the Ray Society.

PROTONINA, <i>Williamson</i> .					
<i>fusiformis</i> , <i>Williamson</i>					
<i>pseudospiralis</i> , <i>Williamson</i>					
ORBULINA, <i>D'Orbigny</i> .					
<i>universa</i> , <i>D'Orb.</i>					
LAGENA, <i>Walker</i> .					
<i>vulgaris</i> , <i>Williamson</i>					
var. <i>clavata</i>					
var. <i>perlucida</i>					
var. <i>semistriata</i>					
var. <i>striata</i>					
var. <i>interrupta</i>					
var. <i>gracilis</i>					
var. <i>substriata</i>					

FORAMINIFERA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
<i>ENTOSOLENIA, Ehrenberg.</i>					
<i>globosa, Walker</i>					
var. <i>lineata</i>					
<i>costata, Williamson</i>					
<i>marginata, Walker</i>					
var. <i>lucida</i>					
var. <i>ornata</i>					
var. <i>lagenoides</i>					
var. <i>quadrata</i>					
<i>squamosa, Mont.</i>					
var. <i>scalariformis</i>					
var. <i>catenulata</i>					
var. <i>hexagona</i>					
<i>LINGULINA, D'Orbigny.</i>					
<i>carinata, D'Orb.</i>					
<i>NODOSARIA, Lamarck.</i>					
<i>radicula, Linn.</i>					
<i>pyrula, D'Orb.</i>					
<i>DENTALINA, D'Orbigny.</i>					
<i>subarcuata, Mont.</i>					
var. <i>jugosa</i>					
<i>legumen, Linn.</i>					
var. <i>linearis</i>					
<i>FRONDICULARIA, DeFrance.</i>					
<i>spathulata, Williamson</i>					
<i>Archiaciana, D'Orb.</i>					
<i>CRISTELLARIA, Lamarck.</i>					
<i>calcar, Linn.</i>					
var. <i>rotifera</i>					
var. <i>oblonga</i>					
<i>subarcuatula, Walker</i>					
var. <i>costata</i>					
<i>NONIONINA, D'Orbigny.</i>					
<i>Barleeana, Williamson</i>					
<i>crassula, Walker</i>					
<i>Jeffreysii, Williamson</i>					
<i>elegans, Williamson</i>					
<i>NUMMULINA, D'Orbigny.</i>					
<i>planulata, Lam.</i>					
<i>POLYSTOMELLA, Lamarck.</i>					
<i>crispa, Linn.</i>					
<i>umbilicatula, Walker</i>					
var. <i>incerta</i>					
<i>PENEROPLIS, Montfort.</i>					
<i>planatus, Ficht. & Moll</i>					
<i>PATELLINA, Williamson.</i>					
<i>corrugata, Williamson</i>					
<i>ROTALINA, D'Orbigny.</i>					
<i>Beccarii, Linn.</i>					

FORAMINIFERA.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
<i>ROTALINA, D'Orbigny.</i>					
<i>inflata, Mont.</i>					
<i>turgida, Williamson.</i>					
<i>oblonga, Williamson</i>					
<i>concamerata, Mont.</i>					
<i>nitida, Williamson</i>					
<i>mamilla, Williamson</i>					
<i>ochracea, Williamson</i>					
<i>fusca, Williamson.</i>					
<i>GLOBIGERINA, D'Orbigny.</i>					
<i>bulloides, D'Orb.</i>					
<i>PLANORBULINA, D'Orbigny.</i>					
<i>vulgaris, D'Orb.</i>					
<i>TRUNCATULINA, D'Orbigny.</i>					
<i>lobatula, Walker</i>					
<i>BULIMINA, D'Orbigny.</i>					
<i>pupoides, D'Orb.</i>					
var. <i>marginata</i>					
var. <i>spinulosa</i>					
var. <i>fusiformis</i>					
var. <i>compressa</i>					
var. <i>convoluta</i>					
<i>elegantissima, D'Orb.</i>					
<i>scabra?, Williamson.</i>					
<i>UVIGERINA, D'Orbigny.</i>					
<i>pygmæa, D'Orb.</i>					
<i>angulosa, Williamson</i>					
<i>CASSIDULINA, D'Orbigny.</i>					
<i>lævigata, D'Orb.</i>					
<i>obtusa, Williamson</i>					
<i>POLYMORPHINA, D'Orbigny.</i>					
<i>lactea, Walker</i>					
var. <i>acuminata</i>					
var. <i>oblonga</i>					
var. <i>fistulosa</i>					
var. <i>concava</i>					
var. <i>communis</i>					
<i>myristiformis, Williamson</i>					
<i>TEXTULARIA, DeFrance.</i>					
<i>euneiformis, D'Orb.</i>					
var. <i>conica</i>					
<i>variabilis, Williamson</i>					
var. <i>spathulata</i>					
var. <i>difformis</i>					
var. <i>lævigata</i>					
<i>BILOCULINA, D'Orbigny.</i>					
<i>ringens, D'Orb.</i>					
var. <i>carinata</i>					

FORAMINIFERA.—SPONGES.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
<i>SPYROLOCULINA, D'Orbigny.</i>					
<i>depressa, D'Orb.</i>					
var. <i>rotundata</i>					
var. <i>cymbium</i>					
<i>MILIOLINA, Williamson.</i>					
<i>trigonula, Lamk.</i>					
<i>seminulum, Linn.</i>					
var. <i>oblonga</i>					
var. <i>disciformis</i>					
<i>bicornis, Walker</i>					
var. <i>elegans</i>					
var. <i>angulata</i>					
<i>VERTEBRALINA, D'Orbigny.</i>					
<i>striata, D'Orb.</i>					
<i>SPIRILLINA, Ehrenberg.</i>					
<i>foliacea, Philippi</i>					
<i>perforata, Schultze</i>					
<i>arenacea, Williamson</i>					
<i>margaritifera, Williamson</i>					

LIST OF BRITISH SPONGES.

<i>TETHYA, Lamarck.</i>					
<i>cranium, Johnst. (not Müll.)</i>					
<i>lyncurium, Johnst.</i>					
<i>GEODIA, Lamarck.</i>					
<i>Zetlandica, Johnst.</i>					
<i>PACHYMATISMA, Bowerbank.</i>					
<i>Johnstonia, Bowb. (HALICHONDRIA, Johnst.)</i>					
<i>HALICHONDRIA, Fleming.</i>					
<i>panicea, Johnst.</i>					
<i>coalita, Johnst.</i>					
<i>coccinea, Bowb. MS.</i>					
<i>glabra, Bowb. MS.</i>					
<i>inconspicua, Bowb. MS.</i>					
<i>caduca, Bowb. MS.</i>					
<i>distorta, Bowb. MS.</i>					
<i>Dickiei, Bowb. MS.</i>					
<i>Batei, Bowb. MS.</i>					
<i>lingua, Bowb. MS.</i>					
<i>corrugata, Bowb. MS.</i>					
<i>granulata, Bowb. MS.</i>					
<i>Thompsoni, Bowb. MS.</i>					
<i>plumosa, Johnst.</i>					

SPONGES.

Species.	Range.	Found living at	Ground.	Frequency	Observations.
	fathoms.	fathoms.			
<i>HALICHONDRIA, Fleming.</i>					
<i>incrustans, Johnst.</i>					
<i>fallax, Bowb. MS.</i>					
<i>paupera, Bowb. MS.</i>					
<i>Clarkei, Bowb. MS.</i>					
<i>variantia, Bowb. MS.</i>					
<i>Hyndmanni, Bowb. MS.</i>					
<i>nigricans, Bowb. MS.</i>					
<i>Ingalli, Bowb. MS.</i>					
<i>fimbriata, Bowb. MS.</i>					
<i>albula, Bowb. MS.</i>					
<i>farinaria, Bowb. MS.</i>					
<i>HYMENIACIDON, Bowerbank (HALICHONDRIA, Johnst.).</i>					
<i>Thomasii, Bowb. MS.</i>					
<i>fragilis, Bowb. MS.</i>					
<i>Brettii, Bowb. MS.</i>					
<i>albescens, Johnst.</i>					
<i>caruncula, Bowb. MS.</i>					
<i>Alderi, Bowb. MS.</i>					
<i>perlævis, Johnst.</i>					
<i>aurea, Johnst.</i>					
<i>pachyderma, Bowb. MS.</i>					
<i>crustula, Bowb. MS.</i>					
<i>sanguinea, Johnst.</i>					
<i>armatura, Bowb. MS.</i>					
<i>floreum, Bowb. MS.</i>					
<i>carnosa, Bowb. MS.</i>					
<i>viridans, Bowb. MS.</i>					
<i>sulphurea, Bowb. MS.</i>					
<i>clavigera, Bowb. MS.</i>					
<i>subclavata, Bowb. MS.</i>					
<i>lactea, Bowb. MS.</i>					
<i>Dujardinii (HALISARCA), Johnst.</i>					
<i>celata, Johnst.</i>					
<i>HALINA, Bowerbank (HALICHONDRIA, Johnst.).</i>					
<i>suberea (HALICHONDRIA), Johnst.</i>					
<i>ficus, Johnst.</i>					
<i>carnosa, Johnst.</i>					
<i>Bucklandi, Bowb. MS.</i>					
<i>ISODICTYA, Bowerbank (HALICHONDRIA, Johnst.).</i>					
<i>Peachii, Bowb. MS.</i>					
<i>rosea, Bowb. MS.</i>					
<i>permollis, Bowb. MS.</i>					
<i>indistincta, Bowb. MS.</i>					
<i>undefinita, Bowb. MS.</i>					
<i>Macandrewii, Bowb. MS.</i>					

SPONGES.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
ISODICTYA, Bowerbank (HALICHONDRIA, Johnst.).					
dichotoma, Bowb. MS.					
cinerea, Johnst.					
ramusculus, Bowb. MS.					
simulo, Bowb. MS.					
mammata, Bowb. MS.					
fucorum, Johnst.					
Alderi, Bowb. MS.					
Normani, Bowb. MS.					
lobata, Johnst.					
Barleci, Bowb. MS.					
gracilis, Bowb. MS.					
Gregorii, Bowb. MS.					
Beauii, Bowb. MS.					
clava, Bowb. MS.					
infundibuliformis, Johnst.					
DESMACIDON, Bowerbank (HALICHONDRIA, Johnst.).					
ægagropila, Johnst.					
fruticosa, Johnst.					
RAPHYRUS, Bowerbank.					
Griffithsii, Bowb. MS.					
DICTYOCYLINDRUS, Bowerbank (HALICHONDRIA, Johnst.).					
stuposus, Johnst.					
Howsei, Bowb. MS.					
ramosus, Johnst.					
aculeatus, Bowb. MS.					
ventilabrum, Bowb. MS.					
fascicularis, Bowb. MS.					
rugosus, Bowb. MS.					
HALICLONA, Bowerbank (HALICHONDRIA, Johnst.).					
palmata, Johnst.					
Montagui, Johnst.					
pygmæa, Bowb.					
seriata, Johnst.					
simulans, Johnst.					
columbæ, Johnst.					
gracilentata, Bowb. MS.					
MICROCIONA, Bowerbank, MS.					
atro-sanguinea, Bowb. MS.					
armata, Bowb. MS.					
carnosa, Bowb. MS.					
ambigua, Bowb. MS.					
lævis, Bowb. MS.					
spinulenta, Bowb. MS.					

SPONGES.

Species.	Range.	Found living at	Ground.	Frequency.	Observations.
	fathoms.	fathoms.			
HYMERAPHIA, <i>Bowerbank, MS.</i>					
<i>vermiculata, Bowb. MS.</i>					
<i>stellifera, Bowb. MS.</i>					
<i>clavata, Bowb. MS.</i>					
HYMEDESMA, <i>Bowerbank, MS.</i>					
<i>Zetlandica, Bowb. MS.</i>					
HALYPHYSEMA, <i>Bowerbank, MS.</i>					
<i>Tumanowiczii, Bowb. MS.</i>					
EUPLECTELLA, <i>Owen.</i>					
<i>mammillaris (HALICHONDRIA),</i>					
<i>Johnst.</i>					
<i>brevis, Bowb. MS.</i>					
<i>robusta, Bowb. MS.</i>					
HALICNEMIA, <i>Bowerbank, MS.</i>					
<i>patera, Bowb. MS.</i>					
PHAKELLIA, <i>Bowb. MS. (HALICHON-</i>					
<i>DRIA, Johnston.)</i>					
<i>ventilabrum, Johnst.</i>					
DYSIDEA, <i>Johnston.</i>					
<i>fragilis, Johnst.</i>					
SPONGIA, <i>Linnaeus.</i>					
<i>pulchella, Johnst.</i>					
<i>limbata, Johnst.</i>					
GRANTIA, <i>Fleming.</i>					
<i>compressa, Johnst.</i>					
<i>ciliata, Johnst.</i>					
<i>ensata, Bowb. MS.</i>					
<i>tessellata, Bowb. MS.</i>					
LEUCONIA, <i>Bowerbank (GRANTIA,</i>					
<i>Johnst.)</i>					
<i>nivea, Johnst.</i>					
<i>fistulosa, Johnst.</i>					
LEUCOSOLENIA, <i>Bowerbank, MS.</i>					
<i>(GRANTIA, Johnst.)</i>					
<i>botryoides, Johnst.</i>					
<i>coriacea, Johnst.</i>					
<i>lacunosa, Johnst.</i>					
<i>contorta, Bowb. MS.</i>					

